

630.7

I26b

no. 669

cop. 8



UNIVERSITY OF
ILLINOIS LIBRARY
AT URBANA-CHAMPAIGN
AGRICULTURE

Illinois
No. 669

CIRCULATION
AGRICULTURE LIBRARY

UNIVERSITY OF ILLINOIS
AGRICULTURE LIBRARY

~~LIBRARY~~

NEW MEXICO STATE
UNIVERSITY

1960 Performance of EXPERIMENTAL CORN HYBRIDS IN ILLINOIS

By Earl R. Leng, R. J. Lambert, M. L. Peasley,
G. L. Ross, and K. E. Williams



Location of
test fields

Bulletin 669

UNIVERSITY OF ILLINOIS · AGRICULTURAL EXPERIMENT STATION

CONTENTS

	Page
MATERIAL TESTED	3
FIELD PROCEDURES AND ANALYSIS OF DATA	5
MEASURING PERFORMANCE	6
TEST RESULTS	6
EXTRÊME NORTHERN ILLINOIS (Woodstock): Double Crosses	7
NORTHERN ILLINOIS (DeKalb): Double Crosses and Three-Way Crosses and Standards	8
WEST NORTH-CENTRAL ILLINOIS (Galesburg): Double Crosses	16
NORTH-CENTRAL ILLINOIS (Peoria): Double Crosses	17
EAST NORTH-CENTRAL ILLINOIS (Ashkum): Double Crosses	18
CENTRAL ILLINOIS (Stanford): Double Crosses	19
WEST-CENTRAL ILLINOIS (Bowen): Double Crosses	20
EAST-CENTRAL ILLINOIS (Urbana): Double Crosses and Three-Way Crosses and Standards	21
WEST SOUTH-CENTRAL ILLINOIS (Greenfield): Double Crosses	29
SOUTHERN ILLINOIS (Brownstown): Double Crosses	30
EXTREME SOUTHERN ILLINOIS (Carbondale): Double Crosses	32
WEST EXTREME SOUTHERN ILLINOIS (Wolf Lake): Double Crosses	33
DOUBLE-CROSS HYBRID NUMBERS, PEDIGREES, AND INDEX TO TABLES	34

Special acknowledgment is due R. W. Jugenheimer, Assistant Dean and Assistant Director of the Agricultural Experiment Station, who developed a considerable amount of the material in this bulletin. Acknowledgment is also due W. C. Jacobs and R. D. Seif for processing the data.

Il 66
no 669
cop. 8

PERFORMANCE OF EXPERIMENTAL CORN HYBRIDS IN ILLINOIS, 1960

By EARL R. LENG, R. J. LAMBERT, M. L. PEASLEY,
G. L. ROSS, and K. E. WILLIAMS¹

ONE OF THE OBJECTIVES OF CORN BREEDERS at the Illinois Agricultural Experiment Station is to develop improved corn inbreds and hybrids for use by seedsmen and farmers of the state. Such development requires considerable breeding work and adequate testing of performance at a number of locations and for a period of years. This bulletin summarizes results of experimental corn hybrid performance trials conducted in 1960. The experimental corn hybrids tested were selected on the basis of their performance in preliminary tests or in advanced tests of previous years.

In 1960 experimental corn hybrids were tested at twelve different locations in the state: Ashkum, Bowen, Brownstown, Carbondale, DeKalb, Galesburg, Greenfield, Peoria, Stanford, Urbana, Wolf Lake, and Woodstock. The maturity series tested at each location, the soil types, the distribution of rainfall during the growing season, dates of planting and harvesting, and planting rates per acre are given in Table 1.

MATERIAL TESTED

A total of 205 corn hybrids, consisting of 109 double crosses, 93 three-way crosses, and 3 single crosses, were tested in advanced corn performance trials in 1960. Most of the hybrids tested were developed by corn breeders at the University of Illinois.

Double crosses tested. Double crosses tested were divided into maturity groups, each consisting of a different set of 25 hybrids. The groups used were based on the AES (Agricultural Experiment Station) maturity series; the groups adapted to Illinois range in maturity from "600" in extreme northern Illinois to the "900" group in southern areas of the state. For testing purposes, hybrids comparable in maturity to those of the AES "800" series were divided into "800" and "850" series. The "800 series" hybrids were grown in north-central Illinois and the "850 series" in central Illinois. Illinois Station hybrids of comparable maturity rating are as follows: 600 series = Illinois 1555A; 700 series = Illinois 3152; 800 series = Illinois 1421; 850 series = Illinois 1570; and 900 series = Illinois 1851.

¹EARL R. LENG, Professor of Agronomy; R. J. LAMBERT, and M. L. PEASLEY, Research Assistants; G. L. ROSS, and K. E. WILLIAMS, Crops Testing Technicians.

Three-way crosses tested. Three-way crosses are useful for evaluating the combining ability of an inbred line. Thirty-one inbreds crossed with three single-cross testers, (WF9 × Oh43), (WF9 × B37), and (B41 × Oh7A), were tested at Brownstown, DeKalb, and Urbana in 1959 and 1960. The test at Brownstown, however, was abandoned in 1960 because of poor stand, so 1959 and 1960 summaries are available only for DeKalb and Urbana.

**Table 1. — GENERAL INFORMATION:
Illinois Experimental Corn Hybrids, 1960**

(All planting rates 16,000 plants per acre, except at Galesburg where it was 18,000, and at Brownstown, where it was 14,000)

Location	Maturity series tested	Soil type	Monthly rainfall (in.)				Date of planting	Date of harvest
			May	June	July	Aug.		
Northern Illinois								
Woodstock	600	Proctor silt loam	5.6	3.9	3.0	2.3	May 15	Oct. 29
DeKalb	600, 700	Flanagan silt loam	6.2	4.1	4.8	3.0	May 24	Nov. 5
North-Central Illinois								
Galesburg	800	Sable silty clay loam	6.1	5.8	2.2	5.4	June 1	Oct. 28
Peoria	700	Muscataine silt loam	6.3	5.4	3.5	5.3	June 8	Nov. 17
Ashkum	700	Milford clay loam	3.1	5.0	1.1	5.1	May 31	Nov. 15
Stanford	800	Muscataine silt loam	3.6	8.3	4.8	2.2	May 12	Oct. 6
Central Illinois								
Bowen	850	Virden silty clay loam	6.7	8.6	3.7	5.0	June 1	Oct. 25
Urbana	850, 900	Brenton silt loam	4.1	6.2	2.8	1.3	May 4	Oct. 6
Southern Illinois								
Greenfield	850, 900	Herrick silt loam	5.8	4.2	3.1	2.1	June 2	Oct. 22
Brownstown	900	Cisne silt loam	5.9	7.2	1.8	2.2	June 9	Nov. 17
Carbondale	900	Weir silt loam	5.5	4.1	1.2	3.8	June 2	Oct. 7
Wolf Lake	900	Riley fine sandy loam	3.9	3.5	2.8	4.6	May 10	Oct. 4

COOPERATORS: EARL HUGHES, *McHenry county*; RALPH ANDERSON, *Knox county*; MELVIN KRAFT, *Iroquois county*; W. T. SCHWENK AND SONS, *Peoria county*; ELDON GOLDEN, *Hancock county*; ROBERT BUTH, *McLean county*; CHARLES ROSS, *Macoupin county*; Shawnee High School, *Union county*. Trials in DeKalb and Champaign counties were located on University of Illinois farms managed by R. E. BELL and C. H. FARNHAM. P. E. JOHNSON, Assistant Professor of Soil Fertility, supervised field operations on the test in Fayette county, and D. R. BROWNING of Southern Illinois University supervised field operations on the Union county and Jackson county test fields.

Performance trials of this type are necessary to properly evaluate improved corn inbreds. The performance of an inbred in a combination with three different single-cross testers is a measure of the combining ability of the inbred line being tested. Tests at a number of locations and for several years more accurately measure combining ability than tests for only one year or at one location.

Availability of material tested. A number of the Illinois Station corn hybrids listed in this report are not yet in commercial production. The Experiment Station release policy is to make available to the public seed of inbred lines that have demonstrated superior performance for desirable agronomic characters. Small amounts of seed (up to 100 kernels) of *released* Illinois inbred lines are available for a nominal fee. Requests for seed of released Illinois inbred lines should be addressed to the Department of Agronomy, University of Illinois, Urbana, Illinois. Station Bulletin 657 lists the Illinois inbred lines released up to and including 1960, and also presents data on some of their important agronomic characteristics. Seed of single crosses that are used as parents for some Illinois Station hybrids reported in this bulletin may be obtained from the Illinois Seed Producers Association, Champaign, Illinois.

FIELD PROCEDURES AND ANALYSIS OF DATA

Method of planting. All test locations except Carbondale were planted with a mounted four-row John Deere tractor planter, slightly modified for planting experimental plots. The Carbondale location was planted by hand. All locations were planted on land prepared in the normal manner for corn. Individual plots were one row 11 hills in length. Planting simulated "power check," with a variable number of kernels being dropped approximately each 20 inches, depending on the planting rate used. All plots were band-treated for weed control with Atrazine at a rate of 12 pounds per acre. The plots were not thinned.

Method of harvest. All plots were harvested with a one-row Ford picker-sheller modified to harvest experimental plots. The shelled corn from each plot was bagged, weighed, and sampled for moisture using a Radson moisture meter. No adjustment was made for dropped ears or for ears on broken stalks that were not harvested.

Field-plot design and analysis of data. The experimental designs used for all trials were lattice designs with 3 replications. All field data were recorded on mark-sense cards and processed with digital computers at the University of Illinois.

MEASURING PERFORMANCE

All hybrids tested were compared for grain yield, kernel moisture, erect plants at harvest, and stand. Data on other agronomic characters such as dropped ears, leaf blight reaction, stalk rot, and smutted plants were recorded when natural conditions permitted measuring true varietal differences.

Yield of grain. Acre yields are reported as shelled corn containing 15.5 percent moisture, the upper limit for No. 2 corn.

Erect plants. A count of erect plants in each plot of an entry was taken at harvest time for each location. Only plants leaning at an angle of 45° or more or broken below the ear were considered lodged; all others were counted as erect.

Stand. A count was made in late summer at all locations of the total number of plants in each plot of a hybrid. The percent stand was computed by comparing the actual number of plants in each plot with the number of kernels planted. Stand differences may have been caused by failure of seed to germinate or by disease, insect damage, cultivation injury, or other factors.

TEST RESULTS

Results from the tests are summarized in Tables 2 to 13. The following facts should be considered when comparing the performance of hybrids in a test.

1. Results covering two and three years at a location are more reliable than results for only one year. The performance of hybrids tested only in 1960 should not be used as a measure of their true ability since further testing will be necessary before valid conclusions can be drawn. This is true of all hybrids tested at Ashkum, Bowen, Carbondale, Galesburg, Greenfield, Stanford, Wolf Lake, and Woodstock. Results from these tests are not ranked by yield but are listed according to hybrid designation. Two- and three-year summaries are available for Brownstown, DeKalb, Peoria, and Urbana, and entries are ranked according to yield in these summaries.

2. Small differences between hybrids do not necessarily indicate that one hybrid is truly superior to another. Interpretation of the data and comparisons between hybrids are made more meaningful by use of certain statistical procedures. One procedure used to compare the difference between hybrids is the "Multiple Range Test."¹ Using this

¹ DUNCAN, D. B. "Multiple Range and Multiple F Tests." *Biometrics* 11 (1) 1-43, 1955.

statistical test, the difference necessary for significance between two or more hybrids can be calculated. Whenever the observed difference between two or more hybrids exceeds the amount calculated for that range, the two hybrids are significantly different. To find the difference necessary for significance the hybrids are first ranked according to performance for a particular character. Then the "number in range" can be computed by counting the hybrids to be compared and the number of hybrids falling between them in performance. For example, if hybrids A and E are to be compared and the rank in performance is A, B, C, D, E, the "number in range" would be 5. When the "number in range" has been determined, the corresponding "difference necessary for significance" can be read from the figures at the bottom of each table. If the observed difference exceeds the "difference necessary for significance," the performances of the hybrids are considered different.

Table 2. — DOUBLE CROSSES OF 600 MATURITY
Tested at Woodstock, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Ill. 1277 (WF9×M14)(L.205×187-2)		92.0	21.7	90.5	95.4
Ill. 1555A (check) (WF9×Oh51A)(L.224×Oh28)		76.5	20.1	98.3	87.8
Ill. 1559B* (WF9×Oh51A)(M14×Oh28)		87.2	22.5	98.3	87.8
Ill. 1861 (WF9×M14)(L.224×Oh28)		68.4	22.5	92.1	87.1
Ill. 1863 (WF9×M14)(L.205×Oh43)		87.2	24.8	92.3	98.4
Ill. 1936 (WF9×Hy2)(M14×B14)		78.0	22.4	93.8	87.8
Ill. 1952 (W64A×A545)(M14×B14)		93.6	22.2	93.4	93.1
Ill. 1955 (W64A×B14)(M14×A297)		83.9	20.9	97.4	83.3
Ill. 1957 (W64A×B14)(M14×A545)		79.9	20.4	98.1	89.3
Ill. 1958 (Oh26A×M14)(B14×A545)		81.5	22.8	94.1	91.6
Ill. 1959 (W64A×M14)(B14×A297)		94.1	23.4	92.8	95.4
Ill. 1960 (W64A×M14)(B14×A545)		86.1	22.7	95.9	96.9
Ill. 1961 (W64A×A239)(B14×A545)		87.5	22.9	100.0	86.3
Ill. 1962 (W64A×A297)(B14×A545)		84.6	21.5	94.5	92.4
Ill. 1969A (WF9×R165)(R168×B14)		80.1	24.4	96.2	81.0
Ill. 3009 (W64A×A297)(B14×B21)		86.0	21.9	95.4	96.2
Ill. 3152 (WF9×M14)(B14×Oh43)		86.8	23.5	97.5	94.6
Ill. 3173 (A545×N24)(B14×Oh43)		84.0	22.8	96.1	90.1
Ill. 3174 (A297×Oh43)(B37×Oh28)		76.3	22.8	92.7	93.9
Ill. 3301 (M14×Oh43)(R168×B14)		85.2	22.8	97.5	96.2
Ill. 3302A-1 ^b (W64A×M14)(R172×B14)		82.5	22.6	94.0	90.1
Ill. 3303 (M14×Oh43)(R172×B14)		81.3	23.0	95.7	90.1
Ill. 3313 (W64A×Oh43)(L12×B14)		69.5	23.8	98.9	78.7
Ill. 6201 (WF9×B14)(R53×Oh7)		77.2	21.2	83.5	96.9
Ill. 6202 (W64A×Oh43)(Oh51×R53)		84.8	20.4	90.6	96.2
Average		83.0	22.5	94.8	91.0
Number in range		Difference necessary for significance			
2		3.9	0.7	1.9	3.0
3-5		4.4	0.8	2.1	3.4
6-10		4.6	0.8	2.3	3.6
11-15		4.7	0.8	2.3	3.7
16-25		4.8	0.9	2.4	3.7

* Illinois Station hybrids with A or B endings in the numerical designation are permutations of a basic arrangement.

^b (-1) indicates that W64A has replaced WF9 in Ill. 3302A (WF9×M14)(R172×B14).

Table 3.—DOUBLE CROSSES OF 600 MATURITY AND 700 MATURITY AND THREE-WAY CROSSES AND STANDARDS
Tested at DeKalb, 1958-1960

Entry	Pedigree	Acre yield	Molsture in grain	Erect plants	Stand
DOUBLE CROSSES OF 600 MATURITY					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Summary: 1958-1960					
III. 3173	(A545×N24)(B14×Oh43)	124.3	28.9	91.0	96.2
III. 3152	(WF9×M14)(B14×Oh43)	122.5	28.2	87.6	96.2
III. 3174	(A297×Oh43)(B37×Oh28)	118.7	27.4	86.9	95.4
III. 1962	(W64A×A297)(B14×A545)	118.3	24.7	86.0	96.3
III. 1936	(WF9×Hy2)(M14×B14)	118.1	28.0	81.9	96.4
III. 1559B	(WF9×Oh51A)(M14×Oh28)	117.9	25.9	78.7	96.6
III. 1952	(W64A×A545)(M14×B14)	116.6	26.7	84.4	96.4
III. 3009	(W64A×A297)(B14×B21)	116.1	25.8	88.3	94.6
III. 1959	(W64A×M14)(B14×A297)	116.1	26.0	90.6	97.1
III. 1961	(W64A×A239)(B14×A545)	114.2	25.1	85.7	97.0
III. 1960	(W64A×M14)(B14×A545)	114.0	26.4	82.7	96.7
III. 1958	(Oh26A×M14)(B14×A545)	113.4	25.0	80.2	97.4
III. 1955	(W64A×B14)(M14×A297)	111.3	24.4	88.1	96.4
III. 1957	(W64A×B14)(M14×A545)	110.6	26.6	80.0	95.1
III. 1555A	(WF9×Oh51A)(I.224×Oh28)	107.5	24.0	77.8	95.5
III. 1277	(WF9×M14)(I.205×187-2)	105.1	26.3	65.7	97.2
Average		115.3	26.2	83.5	96.3
Number in range		Difference necessary for significance			
2	N.S.	2.2	N.S.	N.S.
3-5	N.S.	2.4	N.S.	N.S.
6-10	N.S.	2.5	N.S.	N.S.
11-16	N.S.	2.6	N.S.	N.S.
Summary: 1959-1960					
III. 3303	(M14×Oh43)(R172×B14)	122.7	26.8	92.6	98.1
III. 3301	(M14×Oh43)(R168×B14)	121.7	26.2	97.4	96.9
III. 3173	(A545×N24)(B14×Oh43)	118.1	27.0	92.4	94.3
III. 1962	(W64A×A297)(B14×A545)	117.6	23.5	86.4	96.6
III. 3302A-1	(W64A×M14)(R172×B14)	115.6	25.5	92.6	98.8
III. 3174	(A297×Oh43)(B37×Oh28)	113.3	26.9	87.9	93.5
III. 1960	(W64A×M14)(B14×A545)	113.0	24.7	91.0	95.1
III. 1959	(W64A×M14)(B14×A297)	112.6	24.1	91.7	97.3
III. 3152	(WF9×M14)(B14×Oh43)	112.6	26.5	94.6	94.4
III. 3313	(W64A×Oh43)(L12×B14)	112.5	25.1	95.3	94.8
III. 3009	(W64A×A297)(B14×B21)	112.3	24.7	89.9	93.2
III. 1559B	(WF9×Oh51A)(M14×Oh28)	111.3	25.0	90.2	94.9
III. 1952	(W64A×A545)(M14×B14)	110.3	25.4	90.4	95.9
III. 1936	(WF9×Hy2)(M14×B14)	107.9	25.9	86.8	94.6
III. 1961	(W64A×A239)(B14×A545)	107.4	24.9	87.7	95.5
III. 1969A	(WF9×R165)(R168×B14)	106.9	27.8	95.2	95.1
III. 1958	(Oh26A×M14)(B14×A545)	106.7	23.6	87.3	96.6
III. 1955	(W64A×B14)(M14×A297)	105.8	23.1	86.0	95.0
III. 1957	(W64A×B14)(M14×A545)	105.7	24.4	87.0	93.1
III. 1555A	(WF9×Oh51A)(I.224×Oh28)	100.1	23.3	85.3	94.5
III. 1277	(WF9×M14)(I.205×187-2)	94.7	24.2	83.7	95.9
Average		110.9	25.2	90.1	95.4
Number in range		Difference necessary for significance			
2	N.S.	2.0	N.S.	N.S.
3-5	N.S.	2.2	N.S.	N.S.
6-10	N.S.	2.3	N.S.	N.S.
11-21	N.S.	2.4	N.S.	N.S.

(Table is continued on next page)

Table 3. — DeKalb — continued

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
DOUBLE CROSSES OF 600 MATURITY — concluded					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
1960 results					
III. 1277	(WF9×M14)(I.205×187-2)	106.5	25.4	87.0	93.1
III. 1555A	(check) (WF9×Oh51A)(I.224×Oh28)	104.8	25.5	93.3	92.4
III. 1559B	(WF9×Oh51A)(M14×Oh28)	106.3	26.9	87.2	93.1
III. 1861	(WF9×M14)(I.224×Oh28)	95.5	23.5	77.1	86.3
III. 1863	(WF9×M14)(I.205×Oh43)	108.0	30.0	94.4	93.1
III. 1936	(WF9×Hy2)(M14×B14)	98.9	30.3	89.6	89.3
III. 1952	(W64A×A545)(M14×B14)	98.9	28.5	95.1	93.9
III. 1955	(W64A×B14)(M14×A297)	94.9	25.6	93.4	90.1
III. 1957	(W64A×B14)(M14×A545)	100.1	26.2	92.1	86.3
III. 1958	(Oh26A×M14)(B14×A545)	104.2	24.9	92.7	93.9
III. 1959	(W64A×M14)(B14×A297)	104.8	26.3	96.8	94.6
III. 1960	(W64A×M14)(B14×A545)	110.8	26.8	94.1	90.9
III. 1961	(W64A×A239)(B14×A545)	97.3	27.2	94.2	91.6
III. 1962	(W64A×A297)(B14×A545)	113.4	25.5	95.1	93.9
III. 1969A	(WF9×R165)(R168×B14)	103.1	29.9	96.5	90.9
III. 3009	(W64A×A297)(B14×B21)	103.1	27.2	94.8	87.8
III. 3152	(WF9×M14)(B14×Oh43)	110.0	28.5	96.7	90.1
III. 3173	(A545×N24)(B14×Oh43)	106.9	29.8	95.0	89.3
III. 3174	(A297×Oh43)(B37×Oh28)	107.3	28.5	93.8	87.1
III. 3301	(M14×Oh43)(R168×B14)	110.4	28.2	98.2	93.9
III. 3302A-1	(W64A×M14)(R172×B14)	114.1	26.8	94.6	97.7
III. 3303	(M14×Oh43)(R172×B14)	116.6	28.5	96.0	96.2
III. 3313	(W64A×Oh43)(L12×B14)	106.6	27.2	97.5	91.6
III. 6201	(WF9×B14)(R53×Oh7)	92.6	25.1	82.9	85.6
III. 6202	(W64A×Oh43)(Oh51×R53)	83.7	23.7	86.4	84.0
Average		103.9	27.0	92.5	91.0
Number in range		Difference necessary for significance			
2	4.7	1.0	2.5	N.S.
3-5	5.3	1.1	2.8	N.S.
6-10	5.6	1.1	3.0	N.S.
11-15	5.7	1.2	3.1	N.S.
16-25	5.8	1.2	3.1	N.S.

(Table is continued on next page)

Table 3.—DeKalb—continued

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
DOUBLE CROSSES OF 700 MATURITY					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Summary: 1958-1960					
III. 3152 (WF9×M14)(B14×Oh43)		120.5	28.7	86.4	92.9
III. 1936 (WF9×Hy2)(M14×B14)		119.6	27.0	83.8	93.9
AES 702 (WF9×Hy2)(M14×C103)		105.1	28.3	78.6	97.4
III. 21 (WF9×38-11)(Hy2×187-2)		104.3	27.2	71.9	97.1
III. 1277 (WF9×M14)(I.205×187-2)		102.1	26.8	67.8	97.0
Average		110.3	27.6	77.7	95.7
Number in range		Difference necessary for significance			
2-5		N.S.	N.S.	N.S.	N.S.
Summary: 1959-1960					
III. 3382 (WF9×R109B)(B14×Oh43)		119.1	28.5	93.8	94.3
III. 3381 (WF9×R71)(B14×Oh43)		116.8	28.0	93.5	93.5
III. 3270 (WF9×Oh43)(R74×R168)		115.5	28.1	95.8	97.0
III. 3303 (M14×Oh43)(R172×B14)		114.5	27.2	92.0	92.4
III. 3275 (WF9×Oh43)(R114×R168)		110.6	27.1	95.1	96.3
III. 1936 (WF9×Hy2)(M14×B14)		110.1	24.4	89.7	90.9
III. 3266 (WF9×Oh43)(R74×R109B)		109.8	28.2	89.1	88.1
III. 3152 (WF9×M14)(B14×Oh43)		109.5	27.2	92.8	89.4
III. 3383 (WF9×M14)(R172×Oh43)		109.4	27.2	87.3	94.3
III. 3265 (WF9×Oh43)(R71×R109B)		107.5	28.4	94.8	94.7
III. 21 (WF9×38-11)(Hy2×187-2)		98.7	25.0	83.1	97.0
AES 702 (WF9×Hy2)(M14×C103)		92.0	25.9	89.2	96.2
III. 1277 (WF9×M14)(I.205×187-2)		90.2	24.9	86.8	95.5
Average		108.0	26.9	91.0	93.8
Number in range		Difference necessary for significance			
2-13		N.S.	N.S.	N.S.	N.S.
1960 results					
AES 702 (WF9×Hy2)(M14×C103)		92.1	26.4	85.8	92.4
AES 703 (WF9×Oh43)(B14×B38)		112.0	33.1	97.4	89.3
AES 704 (WF9×Oh43)(B14×B37)		111.4	32.7	98.4	93.9
AES 705 (WF9×B14)(C103×Oh43)		97.2	32.1	97.5	90.9
III. 21 (WF9×38-11)(Hy2×187-2)		112.4	24.9	89.6	94.6
III. 1277 (WF9×M14)(I.205×187-2)		97.5	26.9	93.4	92.4
III. 1922 (WF9×Hy2)(R71×R105)		89.0	33.4	97.2	81.8
III. 1936 (WF9×Hy2)(M14×B14)		103.3	27.3	95.4	81.8
III. 1968 (WF9×B14)(R163×R169)		118.6	27.3	92.9	90.1
III. 1969 (WF9×B14)(R165×R168)		93.0	28.1	93.3	90.9
III. 3022 (WF9×B14)(N22A×Oh43)		99.1	32.4	97.7	96.9
III. 3029 (WF9×B14)(Oh43×Oh45)		95.5	32.8	99.1	93.1
III. 3042 (WF9×B14)(B40×Oh45)		106.6	31.9	95.3	96.2
III. 3152 (check) (WF9×M14)(B14×Oh43)		103.8	30.0	93.2	80.3
III. 3182A (WF9×R105)(R151×R154)		100.6	27.7	94.9	89.3
III. 3265 (WF9×Oh43)(R71×R109B)		110.4	28.4	95.8	90.9
III. 3266 (WF9×Oh43)(R74×R109B)		96.7	30.1	92.2	80.3
III. 3270 (WF9×Oh43)(R74×R168)		106.0	31.1	97.6	94.6
III. 3275 (WF9×Oh43)(R114×R168)		103.2	28.5	98.4	93.9
III. 3303 (M14×Oh43)(R172×B14)		100.1	29.2	94.7	84.8
III. 3315A (WF9×Hy2)(R109B×B14)		85.9	32.6	94.9	93.1
III. 3347 (H49×H55)(R74×R101)		112.9	31.3	92.5	94.6
III. 3381 (WF9×R71)(B14×Oh43)		110.5	31.5	96.3	87.1
III. 3382 (WF9×R109B)(B14×Oh43)		116.8	31.5	94.9	89.3
III. 3383 (WF9×M14)(R172×Oh43)		96.8	29.6	90.7	88.6
Average		102.9	30.0	94.8	90.0
Number in range		Difference necessary for significance			
2		16.4	3.0	6.5	N.S.
3-5		18.2	3.3	7.2	N.S.
6-10		19.3	3.5	7.6	N.S.
11-15		19.7	3.6	7.8	N.S.
16-20		20.0	3.7	7.9	N.S.
21-25		20.1	3.7	7.9	N.S.

(Table is continued on next page)

Table 3. — DeKalb — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (WF9 × Oh43)				
R71.....	99.7	30.1	94.3	88.2
R74.....	98.7	29.1	95.5	81.0
R76.....	101.2	28.8	81.4	83.4
R78.....	99.4	29.4	91.4	88.4
R84.....	92.9	25.8	83.8	83.3
R101.....	96.0	26.4	90.5	88.3
R104.....	99.2	28.0	91.0	81.4
R109B.....	99.9	28.5	96.8	83.0
R112.....	101.8	28.9	96.5	85.2
R113.....	99.3	26.7	90.3	89.5
R114.....	99.4	27.3	95.8	84.3
R134.....	110.9	28.7	93.9	83.1
R135.....	91.7	29.8	87.5	85.6
R151.....	107.6	28.8	90.9	89.1
R154.....	104.1	27.4	90.6	91.3
R158.....	96.7	26.2	92.4	89.5
R159.....	100.2	30.5	93.1	83.3
R166.....	89.7	25.6	88.4	87.6
R168.....	96.0	25.5	96.2	99.5
R172.....	98.5	28.1	92.4	96.8
R180.....	93.6	29.9	88.2	84.2
R181.....	106.2	24.6	86.0	82.0
R182.....	98.8	26.6	97.6	84.5
R183.....	89.2	28.7	89.9	85.5
R192.....	94.7	29.1	93.6	88.6
R193.....	101.0	29.4	94.1	87.3
R194.....	97.9	30.1	88.1	83.6
R195.....	96.0	27.7	91.3	88.2
R196.....	98.4	26.5	97.3	85.4
R197.....	99.7	33.6	91.4	90.0
R198.....	100.2	30.6	92.4	84.1
Average.....	98.7	28.3	91.7	86.6
Inbred lines crossed with (WF9 × B37)				
R71.....	92.3	30.0	96.0	89.2
R74.....	89.5	30.2	97.4	80.7
R76.....	92.3	29.5	89.5	86.1
R78.....	90.9	30.3	85.8	92.5
R84.....	83.6	26.4	81.7	87.0
R101.....	88.1	25.3	87.7	88.2
R104.....	91.1	27.9	81.4	92.3
R109B.....	96.8	29.5	98.4	86.0
R112.....	97.4	29.7	97.5	89.9
R113.....	97.3	28.6	95.4	87.7
R114.....	98.1	30.3	96.4	87.3
R134.....	100.2	30.5	95.6	82.9
R135.....	90.1	30.2	85.5	82.0
R151.....	106.0	30.2	92.4	89.1
R154.....	99.6	27.4	87.9	92.8
R158.....	93.8	28.0	93.9	80.1
R159.....	89.0	30.9	96.6	94.9
R166.....	90.8	28.4	92.1	91.5
R168.....	101.3	26.5	99.6	92.8
R172.....	94.3	28.2	96.9	93.1
R180.....	94.7	30.3	97.5	81.2
R181.....	100.1	26.4	90.5	93.6
R182.....	86.6	26.7	94.7	76.7
R183.....	75.7	29.6	98.1	90.0
R192.....	97.1	28.8	91.0	85.6
R193.....	94.0	29.7	95.8	86.8
R194.....	97.0	31.9	95.7	88.3
R195.....	98.2	27.1	90.9	84.6
R196.....	102.0	29.0	94.2	87.7
R197.....	105.9	32.9	84.6	86.0
R198.....	91.9	32.4	89.5	90.4
Average.....	94.4	29.1	92.6	87.6

(Table is continued on next page)

Table 3. — DeKalb — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960 — continued				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (B41 × Oh7A)				
R71.....	85.8	32.7	83.3	89.1
R74.....	105.0	31.4	92.1	84.1
R76.....	95.6	31.3	73.1	90.4
R78.....	79.9	31.6	72.2	88.1
R84.....	66.5	27.8	79.2	94.1
R101.....	82.9	25.9	83.6	92.7
R104.....	89.6	29.5	83.3	89.5
R109B.....	94.3	33.2	86.9	91.8
R112.....	86.7	32.4	96.0	91.4
R113.....	85.6	28.7	95.8	88.6
R114.....	88.7	30.5	92.0	95.0
R134.....	98.3	33.4	91.2	81.8
R135.....	71.6	34.1	86.3	93.1
R151.....	102.2	28.6	80.4	90.4
R154.....	100.2	29.3	88.9	84.5
R158.....	89.0	28.1	90.7	89.8
R159.....	82.3	33.9	95.7	89.5
R166.....	79.0	30.4	71.5	92.2
R168.....	97.4	27.5	97.1	89.5
R172.....	93.0	29.5	89.5	87.7
R180.....	82.6	31.4	76.8	89.9
R181.....	99.7	25.6	80.4	90.8
R182.....	91.2	29.5	91.6	89.1
R183.....	82.3	30.3	94.3	87.4
R192.....	81.9	30.0	84.6	88.3
R193.....	86.7	30.4	89.6	92.2
R194.....	70.8	33.9	87.5	92.2
R195.....	80.1	27.3	93.7	95.0
R196.....	84.0	30.9	85.4	90.5
R197.....	92.0	31.7	82.1	95.8
R198.....	84.1	33.9	74.5	94.1
Average.....	87.4	30.5	86.1	90.3
Single-cross testers				
WF9 × Oh43.....	100.5	29.6	94.8	85.4
WF9 × B37.....	87.3	31.1	97.6	90.4
B41 × Oh7A.....	65.2	34.1	55.7	92.8
Average.....	84.3	31.6	82.7	89.5
Number in range		Difference necessary for significance		
2.....	N.S.	7.5	8.0	N.S.
3-5.....	N.S.	8.4	8.9	N.S.
6-10.....	N.S.	9.0	9.5	N.S.
11-20.....	N.S.	9.5	10.0	N.S.
21-31.....	N.S.	10.0	10.6	N.S.

(Table is continued on next page)

Table 3. — DeKalb — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960 — continued				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Mean of inbred lines crossed with three testers				
R71.....	92.6	30.9	91.2	88.9
R74.....	97.8	30.3	95.1	82.0
R76.....	96.4	29.9	81.4	86.6
R78.....	90.1	30.5	83.2	89.7
R84.....	81.0	26.7	81.7	88.2
R101.....	89.0	25.9	87.3	89.8
R104.....	93.3	28.5	85.3	87.8
R109B.....	97.1	30.4	94.1	87.0
R112.....	95.3	30.4	96.7	88.9
R113.....	94.1	28.1	93.9	88.6
R114.....	95.5	29.4	94.8	88.9
R134.....	103.2	30.9	93.6	82.6
R135.....	84.5	31.4	86.5	86.9
R151.....	105.3	29.3	88.0	89.5
R154.....	101.4	28.1	89.2	89.6
R158.....	93.2	27.5	92.4	86.5
R159.....	90.6	31.8	95.2	89.3
R166.....	86.6	28.2	84.0	90.5
R168.....	98.2	26.5	97.7	94.0
R172.....	95.3	28.6	93.0	92.6
R180.....	90.4	30.6	87.6	85.2
R181.....	102.0	25.5	85.7	88.8
R182.....	92.2	27.7	94.7	83.5
R183.....	82.4	29.5	94.1	87.7
R192.....	91.3	29.3	89.8	87.6
R193.....	93.9	29.9	93.2	88.8
R194.....	88.6	32.0	90.5	88.1
R195.....	91.5	27.4	92.0	89.3
R196.....	94.8	28.8	92.3	87.9
R197.....	99.3	32.8	86.1	90.7
R198.....	92.1	32.3	85.5	89.6
Average.....	93.5	29.3	90.2	88.2
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS				
Inbred lines crossed with (WF9 × Oh43)				
R71.....	90.5	31.7	92.6	81.7
R74.....	98.5	31.0	97.9	70.4
R76.....	94.7	30.5	89.4	75.7
R78.....	88.3	31.2	92.8	84.8
R84.....	91.5	26.2	84.9	80.2
R101.....	89.9	27.1	89.1	82.5
R104.....	98.5	29.7	98.0	70.4
R109B.....	94.2	30.0	97.0	76.4
R112.....	93.1	31.5	97.0	78.0
R113.....	89.9	28.5	95.9	82.5
R114.....	93.7	29.8	98.8	77.2
R134.....	93.7	30.1	94.1	77.2
R135.....	88.9	33.3	95.5	84.0
R151.....	89.9	30.4	96.3	82.5
R154.....	84.1	29.5	93.6	90.8
R158.....	89.9	27.5	92.6	82.5
R159.....	95.8	31.8	96.9	74.2
R166.....	88.3	25.4	89.2	84.8
R168.....	77.7	27.3	99.2	99.9
R172.....	80.9	30.0	91.4	95.4
R180.....	94.2	32.2	89.3	76.4
R181.....	96.9	27.5	94.8	72.7
R182.....	92.1	27.6	99.0	79.5
R183.....	93.7	30.5	99.1	77.2
R192.....	90.5	30.8	96.1	81.7
R193.....	91.5	31.5	95.0	80.2
R194.....	96.3	31.4	95.7	73.4
R195.....	91.0	30.0	93.0	81.0
R196.....	94.7	27.3	98.1	75.7
R197.....	88.3	36.5	95.8	84.8
R198.....	95.3	32.8	96.9	74.9
Average.....	91.5	30.0	94.7	80.3

(Table is continued on next page)

Table 3. — DeKalb — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS — continued				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (WF9 × B37)				
R71.....	85.1	30.4	94.1	89.3
R74.....	85.7	31.4	95.7	88.6
R76.....	92.1	31.4	89.4	79.5
R78.....	84.1	31.6	86.9	90.8
R84.....	91.0	27.3	77.1	81.0
R101.....	90.5	25.2	87.0	81.7
R104.....	86.2	28.9	83.0	87.8
R109B.....	92.6	31.1	98.0	78.7
R112.....	89.4	32.4	97.2	83.3
R113.....	92.1	31.7	97.0	79.5
R114.....	92.1	32.1	94.7	79.5
R134.....	95.3	32.3	97.0	74.9
R135.....	88.9	33.5	87.0	84.0
R151.....	89.9	31.0	93.4	82.5
R154.....	84.6	28.6	94.2	90.1
R158.....	95.8	29.8	96.7	74.2
R159.....	83.5	33.2	98.4	91.6
R166.....	86.2	29.8	93.8	87.8
R168.....	85.1	27.5	100.0	89.3
R172.....	85.7	29.3	98.2	88.6
R180.....	96.9	32.3	98.1	72.7
R181.....	85.1	28.7	91.5	89.3
R182.....	89.9	28.5	93.2	82.5
R183.....	88.3	30.1	98.2	84.8
R192.....	92.6	29.8	93.4	78.7
R193.....	93.1	31.0	99.0	78.0
R194.....	89.4	33.2	95.7	83.3
R195.....	94.7	29.0	91.0	75.7
R196.....	91.5	30.8	94.4	80.2
R197.....	93.1	34.1	89.1	78.0
R198.....	88.9	33.4	94.5	84.0
Average.....	89.7	30.6	93.4	82.9
Inbred lines crossed with (B41 × Oh7A)				
R71.....	89.5	33.8	79.7	83.3
R74.....	95.9	33.4	92.3	74.2
R76.....	88.9	32.3	71.2	84.0
R78.....	91.6	32.1	81.0	80.2
R84.....	83.6	27.0	77.6	91.6
R101.....	85.7	25.9	86.1	88.6
R104.....	90.0	30.9	91.5	82.5
R109B.....	87.3	35.2	82.9	86.3
R112.....	84.1	35.0	96.8	86.3
R113.....	90.5	31.2	97.2	81.7
R114.....	82.5	32.5	93.6	93.1
R134.....	99.1	35.6	95.3	69.6
R135.....	85.7	37.6	82.5	88.6
R151.....	88.9	29.1	84.0	84.0
R154.....	95.9	31.2	94.9	74.2
R158.....	86.8	29.0	89.5	87.0
R159.....	90.0	34.8	96.2	82.5
R166.....	86.8	31.6	79.2	87.0
R168.....	90.0	28.5	97.2	82.5
R172.....	91.6	31.8	92.0	80.2
R180.....	89.5	33.6	81.4	83.3
R181.....	88.4	27.5	85.3	84.8
R182.....	89.5	30.9	90.1	83.3
R183.....	90.5	31.6	93.9	81.7
R192.....	90.0	30.6	84.6	82.5
R193.....	86.8	32.6	92.0	87.0
R194.....	86.8	33.8	84.6	87.0
R195.....	83.1	27.2	91.0	92.3
R196.....	87.9	33.0	86.6	85.5
R197.....	82.5	32.6	80.9	93.1
R198.....	84.1	36.0	84.5	90.8
Average.....	88.5	31.9	87.6	84.5

(Table is concluded on next page)

Table 3. — DeKalb — concluded

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS — concluded					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Single-cross testers					
WF9×Oh43.....		97.0	31.3	94.7	72.7
WF9×B37.....		90.6	34.2	98.2	81.8
B41×Oh7A.....		85.3	34.2	64.4	88.6
Average.....		91.0	33.2	85.8	81.1
Standards					
III. 1851 (C103×38-11)(Oh7×Cl.21E).....		85.8	32.6	94.0	88.6
III. 3049 (Hy2×WF9)(R71×R109B).....		80.5	32.3	90.5	95.4
III. 3152A (M14×B14)(WF9×Oh43).....		90.6	30.0	98.2	81.8
III. 3347 (R74×R101)(H49×H55).....		83.1	34.8	91.6	93.2
Average.....		85.0	32.4	93.6	89.8
Number in range		Difference necessary for significance			
2.....		N.S.	10.9	9.9	N.S.
3-5.....		N.S.	12.2	11.1	N.S.
6-10.....		N.S.	13.0	11.8	N.S.
11-20.....		N.S.	13.7	12.4	N.S.
21-31.....		N.S.	13.9	12.6	N.S.
Mean of inbred lines crossed with three testers					
R71.....		88.4	32.0	88.8	84.8
R74.....		93.4	32.0	95.4	77.7
R76.....		92.0	31.5	83.4	79.8
R78.....		88.0	31.6	87.0	85.3
R84.....		88.8	26.8	79.9	84.3
R101.....		88.7	26.1	87.5	84.3
R104.....		91.6	29.9	90.9	80.3
R109B.....		91.4	32.1	92.6	80.5
R112.....		88.9	33.0	97.0	82.5
R113.....		90.9	30.5	96.8	81.3
R114.....		89.4	31.5	95.8	83.3
R134.....		96.0	32.8	95.5	74.0
R135.....		87.8	34.9	88.4	85.6
R151.....		89.6	30.2	91.3	83.0
R154.....		88.2	29.8	94.3	85.1
R158.....		90.9	28.8	93.0	81.3
R159.....		89.8	33.3	97.2	82.8
R166.....		87.2	29.0	87.4	86.6
R168.....		84.3	27.8	98.8	90.6
R172.....		86.1	30.4	93.9	88.1
R180.....		93.6	32.7	89.6	77.5
R181.....		90.2	27.9	90.6	82.3
R182.....		90.5	29.0	94.2	81.8
R183.....		90.9	30.8	97.1	81.3
R192.....		91.1	30.4	91.5	81.0
R193.....		90.5	31.7	95.4	81.8
R194.....		90.9	32.8	92.1	81.3
R195.....		89.7	28.8	91.7	83.0
R196.....		91.4	30.4	93.1	80.5
R197.....		88.0	34.4	88.7	85.3
R198.....		89.4	34.1	92.0	83.3
Average.....		89.9	30.9	92.0	82.6

Table 4. — DOUBLE CROSSES OF 800 MATURITY
Tested at Galesburg, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 705	(WF9×B14)(C103×Oh43)	94.8	24.6	97.8	92.0
AES 809	(WF9×P8)(C103×Oh43)	95.6	27.4	85.7	84.0
AES 810	(WF9×H50)(Oh7B×Oh45)	106.8	27.2	95.7	86.0
III. 1421	(check) (WF9×Hy2)(P8×Oh7)	111.9	27.3	87.0	89.3
III. 1983	(WF9×38-11)(Hy2×B14)	105.2	24.0	90.0	86.6
III. 1996	(Hy2×Oh7)(B14×C103)	107.2	26.0	92.7	90.6
III. 3042	(WF9×B14)(B40×Oh45)	108.8	26.2	89.7	96.6
III. 3049	(WF9×Hy2)(R71×R109B)	91.5	26.7	90.5	85.3
III. 3080	(WF9×Hy2)(R101×Oh451)	94.1	25.2	74.8	90.0
III. 3160	(WF9×Oh7)(B14×Oh43)	115.6	26.3	96.8	84.6
III. 3183	(WF9×R154)(R105×R153)	100.3	28.1	87.5	88.6
III. 3237	(WF9×R101)(R151×R154)	115.0	24.7	92.9	81.3
III. 3244	(WF9×R151)(R105×R153)	104.1	26.6	87.9	93.3
III. 3291	(WF9×P8)(B14×Oh43)	108.1	25.3	96.5	92.6
III. 3294	(WF9×P8)(Hy2×C103)	104.1	28.0	89.6	96.6
III. 3343	(H49×H55)(R71×R74)	118.4	28.9	90.4	98.0
III. 3346	(H49×H55)(R71×R168)	130.2	27.9	97.1	80.0
III. 3348	(H49×H55)(R74×R109B)	125.4	26.7	89.6	89.3
III. 3351	(H49×H55)(R109B×R168)	108.6	26.6	91.4	93.3
III. 3384	(WF9×Oh41)(Hy2×Oh7)	93.4	24.3	78.0	91.3
III. 8001	(Hy2×R138)(Oh7×Oh7B)	114.1	28.0	84.6	88.0
III. 8002 ^a	[(Hy2×B14)Hy2][(Oh7×C103)Oh7]	119.6	26.1	89.4	94.0
III. 8003	(WF9×Oh7)(H55×C103)	120.6	26.2	93.4	90.0
III. 8004	(WF9×Hy2)(R74×B14)	93.0	27.1	93.0	86.0
U.S. 13	(WF9×38-11)(Hy2×L317)	119.9	27.6	89.8	88.0
Average		108.3	26.5	90.1	89.4
Number in range		Difference necessary for significance			
	2	16.1	N.S.	8.5	N.S.
	3-5	17.8	N.S.	9.5	N.S.
	6-10	18.9	N.S.	10.0	N.S.
	11-15	19.4	N.S.	10.3	N.S.
	16-20	19.6	N.S.	10.4	N.S.
	21-25	19.7	N.S.	10.5	N.S.

^a Back-cross hybrid.

Table 5. — DOUBLE CROSSES OF 700 MATURITY
Tested at Peoria, 1958-1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
Summary: 1958-1960					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
III. 3022 (WF9×B14)(N22A×Oh43)		107.3	22.0	93.5	93.1
III. 3029 (WF9×B14)(Oh43×Oh45)		103.8	22.7	93.4	91.7
III. 3042 (WF9×B14)(B40×Oh45)		102.0	24.0	90.2	93.1
III. 1968 (WF9×B14)(R163×R169)		100.1	20.9	87.9	93.7
AES 703 (WF9×Oh43)(B14×B38)		99.2	21.7	94.9	92.3
III. 1969 (WF9×B14)(R165×R168)		98.6	20.7	90.8	95.3
AES 705 (WF9×B14)(C103×Oh43)		96.8	23.0	90.3	94.6
III. 21 (WF9×38-11)(Hy2×187-2)		95.7	20.4	76.7	93.3
AES 704 (WF9×Oh43)(B14×B37)		93.1	21.9	97.8	90.4
AES 702 (WF9×Hy2)(M14×C103)		92.3	21.4	87.5	90.6
Average		98.9	21.9	90.3	92.8
Number in range		Difference necessary for significance			
2-10		N.S.	N.S.	N.S.	N.S.
Summary: 1959-1960					
III. 3347 (H49×H55)(R74×R101)		98.3	25.2	86.4	89.6
III. 3022 (WF9×B14)(N22A×Oh43)		95.3	23.9	95.6	91.3
III. 3182A (WF9×R105)(R151×R154)		90.5	24.3	75.0	93.7
III. 3029 (WF9×B14)(Oh43×Oh45)		87.8	24.2	97.0	87.6
AES 703 (WF9×Oh43)(B14×B38)		87.1	23.9	94.8	90.5
III. 1968 (WF9×B14)(R163×R169)		86.8	22.9	87.2	90.6
III. 3042 (WF9×B14)(B40×Oh45)		86.7	25.7	91.5	90.0
III. 1969 (WF9×B14)(R165×R168)		85.4	22.3	90.3	93.0
III. 21 (WF9×38-11)(Hy2×187-2)		83.0	21.6	85.2	92.1
AES 702 (WF9×Hy2)(M14×C103)		81.2	22.5	92.6	86.7
III. 3315A (WF9×Hy2)(R109B×B14)		79.9	22.5	94.1	89.0
AES 705 (WF9×B14)(C103×Oh43)		79.7	25.6	93.0	91.9
AES 704 (WF9×Oh43)(B14×B37)		75.4	23.5	97.6	86.4
Average		85.9	23.7	90.8	90.2
Number in range		Difference necessary for significance			
2		N.S.	2.4	9.8	N.S.
3-5		N.S.	2.6	10.7	N.S.
6-13		N.S.	2.7	11.0	N.S.
1960 results					
AES 702 (WF9×Hy2)(M14×C103)		64.6	23.5	92.4	79.5
AES 703 (WF9×Oh43)(B14×B38)		72.8	26.0	95.8	87.1
AES 704 (WF9×Oh43)(B14×B37)		65.5	24.8	97.3	80.3
AES 705 (WF9×B14)(C103×Oh43)		66.9	28.9	88.0	88.6
III. 21 (WF9×38-11)(Hy2×187-2)		77.7	22.4	82.2	90.9
III. 1277 (WF9×M14)(L205×187-2)		63.5	23.2	89.7	81.0
III. 1922 (WF9×Hy2)(R71×R105)		78.8	26.6	95.6	87.8
III. 1936 (WF9×Hy2)(M14×B14)		47.4	23.8	86.8	97.7
III. 1968 (WF9×B14)(R163×R169)		64.4	25.5	82.8	87.8
III. 1969 (WF9×B14)(R165×R168)		66.6	23.9	87.4	90.1
III. 3022 (WF9×B14)(N22A×Oh43)		74.4	26.2	95.6	89.3
III. 3029 (WF9×B14)(Oh43×Oh45)		67.8	26.1	96.3	83.3
III. 3042 (WF9×B14)(B40×Oh45)		59.1	27.6	86.6	84.8
III. 3152 (check) (WF9×M14)(B14×Oh43)		67.5	23.5	94.4	83.3
III. 3182A (WF9×R105)(R151×R154)		63.2	25.1	81.7	90.9
III. 3265 (WF9×Oh43)(R71×R109B)		70.1	26.6	94.6	87.8
III. 3266 (WF9×Oh43)(R74×R109B)		68.3	28.0	92.9	86.3
III. 3270 (WF9×Oh43)(R74×R168)		69.2	24.1	95.7	84.8
III. 3275 (WF9×Oh43)(R114×R168)		75.3	23.3	97.5	83.3
III. 3303 (M14×Oh43)(R172×B14)		84.1	24.1	94.9	88.6
III. 3315A (WF9×Hy2)(R109B×B14)		65.7	23.2	94.6	83.3
III. 3347 (H49×H55)(R74×R101)		84.2	26.8	83.2	83.0
III. 3381 (WF9×R71)(B14×Oh43)		61.4	25.4	89.7	84.3
III. 3382 (WF9×R109B)(B14×Oh43)		73.4	27.1	94.4	79.5
III. 3383 (WF9×M14)(R172×Oh43)		74.5	24.4	98.2	90.9
Average		69.1	25.2	91.5	86.2
Number in range		Difference necessary for significance			
2		15.9	3.2	11.3	N.S.
3-5		17.6	3.6	12.5	N.S.
6-10		18.7	3.8	13.2	N.S.
11-15		19.1	3.9	13.6	N.S.
16-20		19.4	3.9	13.7	N.S.
21-25		19.5	3.9	13.8	N.S.

Table 6. — DOUBLE CROSSES OF 700 MATURITY
Tested at Ashkum, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 702	(WF9×Hy2)(M14×C103)	62.6	21.4	88.0	95.4
AES 703	(WF9×Oh43)(B14×B38)	76.1	25.6	95.5	90.1
AES 704	(WF9×Oh43)(B14×B37)	83.9	24.3	95.0	91.6
AES 705	(WF9×B14)(C103×Oh43)	80.1	24.1	96.0	94.6
III. 21	(WF9×38-11)(Hy2×187-2)	77.7	21.9	92.9	97.7
III. 1277	(WF9×M14)(I.205×187-2)	81.4	22.0	90.6	88.6
III. 1922	(WF9×Hy2)(R71×R105)	71.7	26.5	95.2	90.9
III. 1936	(WF9×Hy2)(M14×B14)	81.5	24.0	96.1	97.7
III. 1968	(WF9×B14)(R163×R169)	85.2	19.1	89.5	93.1
III. 1969	(WF9×B14)(R165×R168)	87.8	20.3	89.2	96.2
III. 3022	(WF9×B14)(N22A×Oh43)	80.7	23.6	97.7	95.4
III. 3029	(WF9×B14)(Oh43×Oh45)	84.4	21.1	96.4	86.3
III. 3042	(WF9×B14)(B40×Oh45)	83.6	24.2	91.2	97.7
III. 3152	(check) (WF9×M14)(B14×Oh43)	84.0	21.5	91.7	93.1
III. 3182A	(WF9×R105)(R151×R154)	80.8	23.3	89.4	92.4
III. 3265	(WF9×Oh43)(R71×R109B)	85.9	25.8	97.7	96.9
III. 3266	(WF9×Oh43)(R74×R109B)	89.2	23.7	96.0	99.2
III. 3270	(WF9×Oh43)(R74×R168)	89.7	24.0	92.6	96.2
III. 3275	(WF9×Oh43)(R114×R168)	77.8	22.7	91.5	98.4
III. 3303	(M14×Oh43)(R172×B14)	81.9	22.2	94.5	97.7
III. 3315A	(WF9×Hy2)(R109B×B14)	74.3	24.4	94.3	92.4
III. 3347	(H49×H55)(R74×R101)	89.1	25.0	91.2	95.4
III. 3381	(WF9×R71)(B14×Oh43)	82.3	24.3	93.1	99.2
III. 3382	(WF9×R109B)(B14×Oh43)	88.7	22.1	94.6	98.4
III. 3383	(WF9×M14)(R172×Oh43)	87.3	22.4	93.6	95.4
Average		81.9	23.2	93.3	94.8
Number in range		Difference necessary for significance			
	2	N.S.	3.4	N.S.	N.S.
	3-5	N.S.	3.8	N.S.	N.S.
	6-10	N.S.	4.0	N.S.	N.S.
	11-15	N.S.	4.1	N.S.	N.S.
	16-25	N.S.	4.2	N.S.	N.S.

Table 7.—DOUBLE CROSSES OF 800 MATURITY
Tested at Stanford, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 705 (WF9×B14)(C103×Oh43)		105.7	21.8	96.8	90.1
AES 809 (WF9×P8)(C103×Oh43)		97.5	21.2	97.4	93.1
AES 810 (WF9×H50)(Oh7B×Oh45)		105.2	19.5	98.4	87.8
Ill. 1421 (check) (WF9×Hy2)(P8×Oh7)		94.7	19.0	98.3	93.9
Ill. 1983 (WF9×38-11)(Hy2×B14)		116.2	17.6	96.9	100.0
Ill. 1996 (Hy2×Oh7)(B14×C103)		105.8	20.4	96.7	90.9
Ill. 3042 (WF9×B14)(B40×Oh45)		108.0	20.7	97.6	93.9
Ill. 3049 (WF9×Hy2)(R71×R109B)		90.4	20.1	98.2	90.1
Ill. 3080 (WF9×Hy2)(R101×Oh451)		104.8	21.3	93.8	96.9
Ill. 3160 (WF9×Oh7)(B14×Oh43)		113.9	18.0	99.1	88.6
Ill. 3183 (WF9×R154)(R105×R153)		92.0	20.8	100.0	83.3
Ill. 3237 (WF9×R101)(R151×R154)		102.0	16.4	96.6	90.9
Ill. 3244 (WF9×R151)(R105×R153)		86.4	20.7	100.0	92.4
Ill. 3291 (WF9×P8)(B14×Oh43)		119.0	21.3	96.2	100.0
Ill. 3294 (WF9×P8)(Hy2×C103)		99.0	20.0	92.8	96.2
Ill. 3343 (H49×H55)(R71×R74)		110.2	21.5	98.2	90.1
Ill. 3346 (H49×H55)(R71×R168)		121.6	20.8	99.1	92.4
Ill. 3348 (H49×H55)(R74×R109B)		117.2	22.3	99.2	96.2
Ill. 3351 (H49×H55)(R109B×R168)		108.3	22.2	96.8	95.4
Ill. 3384 (WF9×Oh41)(Hy2×Oh7)		99.9	19.1	96.6	89.3
Ill. 8001 (Hy2×R138)(Oh7×Oh7B)		110.8	20.3	96.0	93.1
Ill. 8002 [(Hy2×B14)Hy2][(Oh7×C103)Oh7]		108.6	19.2	98.3	95.4
Ill. 8003 (WF9×Oh7)(H55×C103)		113.2	19.0	97.5	92.4
Ill. 8004 (WF9×Hy2)(R74×B14)		96.7	20.4	99.1	92.4
U.S. 13 (WF9×38-11)(Hy2×L317)		97.0	18.5	95.3	99.2
Average		105.0	20.1	97.4	93.0
Number in range		Difference necessary for significance			
2		N.S.	2.0	N.S.	N.S.
3-5		N.S.	2.3	N.S.	N.S.
6-10		N.S.	2.4	N.S.	N.S.
11-25		N.S.	2.5	N.S.	N.S.

Table 8. — DOUBLE CROSSES OF 850 MATURITY
Tested at Bowen, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 805	(WF9×38-11)(C103×Oh45)	80.3	21.5	79.9	90.9
III. 1332	(WF9×38-11)(Hy2×Oh7)	104.2	20.8	87.8	91.6
III. 1570	(check) (WF9×38-11)(Hy2×Oh41)	94.0	20.9	84.5	93.9
III. 1660	(Oh7×C1.21E)(K4×K201)	115.4	27.8	78.5	87.8
III. 1976	(Oh7×C1.21E)(38-11×Oh41)	94.5	23.3	88.0	88.6
III. 1978	(WF9×Oh7A)(C103×38-11)	87.6	21.7	89.0	75.0
III. 1996	(Hy2×Oh7)(C103×B14)	104.2	21.6	88.9	88.6
III. 3154	(K201×C1.21E)(R132×R134)	94.1	27.8	86.9	75.7
III. 3190	(K201×C103)(Ky126×Oh7B)	109.4	24.1	81.8	90.9
III. 3344	(H49×H55)(R71×R105)	106.2	25.2	93.9	86.3
III. 3347	(H49×H55)(R74×R101)	104.5	22.1	89.3	86.3
III. 3348	(H49×H55)(R74×R109B)	102.5	24.8	96.7	93.9
III. 3350	(H49×H55)(R101×Oh41)	91.6	23.3	95.2	94.6
III. 3351	(H49×H55)(R109B×R168)	102.4	22.7	84.8	88.6
III. 3354	(H49×H51)(R71×R105)	91.5	24.1	84.1	94.6
III. 3357	(H49×H51)(R74×R101)	94.7	22.9	94.7	90.1
III. 3367	(WF9×R74)(Oh7×C1.21E)	109.1	22.9	92.1	87.1
III. 3373	(WF9×C103)(R101×Oh41)	83.9	21.7	88.5	86.3
III. 6021	(R75×R76)(R84×K4)	91.7	20.6	86.8	92.4
III. 6052	(R78×38-11)(R84×K4)	89.1	21.9	74.5	87.8
III. 8005	(H49×H55)(B14×C103)	93.8	22.9	91.4	87.1
Ind. 851	(H49×H55)(H59×B14)	103.3	24.9	86.5	86.3
Ind. 873	(H49×H52)(H59×B14)	98.3	24.0	93.7	87.8
Ind. 874	(H49×H52)(H59×H60)	88.9	22.3	90.3	95.4
U.S. 13	(WF9×38-11)(Hy2×L317)	67.0	21.1	78.3	90.9
Average		96.1	23.1	87.4	88.7
Number in range		Difference necessary for significance			
2		N.S.	1.7	10.5	9.2
3-5		N.S.	1.9	11.6	10.2
6-10		N.S.	2.0	12.3	10.8
11-15		N.S.	2.0	12.6	11.1
16-20		N.S.	2.1	12.7	11.2
21-25		N.S.	2.1	12.8	11.3

Table 9.—DOUBLE CROSSES OF 850 MATURITY AND 900 MATURITY AND THREE-WAY CROSSES AND STANDARDS
Tested at Urbana, 1958-1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
DOUBLE CROSSES OF 850 MATURITY					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Summary: 1958-1960					
U.S. 13 (WF9×38-11)(Hy2×L317)		97.4	21.4	89.0	95.6
Ill. 6052 (R78×38-11)(R84×K4)		95.9	21.9	85.7	94.2
Ill. 1570 (WF9×38-11)(Hy2×Oh41)		95.1	21.2	93.4	93.7
Ill. 6021 (R75×R76)(R84×K4)		92.9	20.9	91.0	96.2
Ill. 1976 (Oh7×Cl.21E)(38-11×Oh41)		92.4	20.9	93.3	94.1
Ill. 1978 (WF9×Oh7A)(C103×38-11)		90.9	22.0	90.7	92.5
Ill. 1332 (WF9×38-11)(Hy2×Oh7)		90.7	20.3	97.7	95.8
Ill. 1996 (Hy2×Oh7)(C103×B14)		88.6	20.2	96.3	90.3
AES 805 (WF9×38-11)(C103×Oh45)		81.1	21.7	94.7	93.1
Average		91.7	21.2	92.4	93.9
Number in range		Difference necessary for significance			
2-9		N.S.	N.S.	N.S.	N.S.
Summary: 1959-1960					
Ill. 3350 (H49×H55)(R101×Oh41)		102.9	23.8	92.9	96.2
Ill. 3347 (H49×H55)(R74×R101)		97.9	22.4	96.1	88.6
Ill. 3357 (H49×H51)(R74×R101)		92.1	23.4	95.8	91.5
Ill. 3354 (H49×H51)(R71×R105)		91.8	25.1	87.6	92.1
Ill. 3344 (H49×H55)(R71×R105)		90.8	24.6	95.9	92.9
Ill. 3367 (WF9×R74)(Oh7×Cl.21E)		89.2	21.8	95.3	90.5
Ill. 3351 (H49×H55)(R109B×R168)		87.0	23.3	94.0	95.8
Ill. 6052 (R78×38-11)(R84×K4)		85.7	23.7	91.3	96.3
U.S. 13 (WF9×38-11)(Hy2×L317)		84.4	22.7	94.2	95.1
Ill. 6021 (R75×R76)(R84×K4)		83.2	22.2	95.5	95.2
Ill. 1976 (Oh7×Cl.21E)(38-11×Oh41)		82.5	21.4	96.2	92.9
Ill. 1570 (WF9×38-11)(Hy2×Oh41)		82.1	22.2	95.8	93.6
Ill. 3373 (WF9×C103)(R101×Oh41)		80.7	21.3	95.2	93.2
Ill. 3348 (H49×H55)(R74×R109B)		80.7	24.7	93.2	90.5
Ill. 1332 (WF9×38-11)(Hy2×Oh7)		76.6	21.1	99.1	95.0
Ill. 1978 (WF9×Oh7A)(C103×38-11)		74.1	23.6	98.9	89.1
Ill. 1996 (Hy2×Oh7)(C103×B14)		73.5	21.0	95.9	87.9
AES 805 (WF9×38-11)(C103×Oh45)		63.6	22.8	97.3	91.7
Average		84.4	22.8	95.0	92.7
Number in range		Difference necessary for significance			
2-18		N.S.	N.S.	N.S.	N.S.
1960 results					
AES 805 (WF9×38-11)(C103×Oh45)		77.6	27.4	98.1	84.8
Ill. 1332 (WF9×38-11)(Hy2×Oh7)		93.9	24.7	99.0	90.1
Ill. 1570 (check) (WF9×38-11)(Hy2×Oh41)		89.9	26.2	97.1	87.8
Ill. 1660 (Oh7×Cl.21E)(K4×K201)		76.5	22.8	100.0	83.3
Ill. 1976 (Oh7×Cl.21E)(38-11×Oh41)		79.0	23.7	100.0	87.8
Ill. 1978 (WF9×Oh7A)(C103×38-11)		95.6	27.1	100.0	81.0
Ill. 1996 (Hy2×Oh7)(C103×B14)		72.6	24.0	100.0	77.2
Ill. 3154 (K201×Cl.21E)(R132×R134)		106.3	27.7	98.3	87.8
Ill. 3190 (K201×C103)(Ky126×Oh7B)		82.1	28.1	100.0	79.5
Ill. 3344 (H49×H55)(R71×R105)		92.7	25.4	96.7	88.6
Ill. 3347 (H49×H55)(R74×R101)		94.0	26.4	100.0	82.5
Ill. 3348 (H49×H55)(R74×R109B)		71.8	28.3	94.5	81.8
Ill. 3350 (H49×H55)(R101×Oh41)		103.8	25.3	99.2	93.1
Ill. 3351 (H49×H55)(R109B×R168)		93.1	26.4	100.0	91.6
Ill. 3354 (H49×H51)(R71×R105)		92.5	26.8	99.0	85.6
Ill. 3357 (H49×H51)(R74×R101)		85.8	26.2	100.0	86.3
Ill. 3367 (WF9×R74)(Oh7×Cl.21E)		92.6	24.1	100.0	81.8
Ill. 3373 (WF9×C103)(R101×Oh41)		79.4	24.5	100.0	87.8
Ill. 6021 (R75×R76)(R84×K4)		106.4	25.0	100.0	92.4
Ill. 6052 (R78×38-11)(R84×K4)		108.1	27.0	100.0	100.0
Ill. 8005 (H49×H55)(B14×C103)		86.7	28.0	100.0	89.3
Ind. 851 (H49×H55)(H59×B14)		95.2	23.8	99.1	93.1
Ind. 873 (H49×H52)(H59×B14)		101.0	24.1	100.0	94.6
Ind. 874 (H49×H52)(H59×H60)		84.5	24.8	98.2	89.3
U.S. 13 (WF9×38-11)(Hy2×L317)		99.7	26.7	100.0	90.9
Average		90.4	25.8	99.2	87.5
Number in range		Difference necessary for significance			
2-25		N.S.	N.S.	N.S.	N.S.

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
DOUBLE CROSSES OF 900 MATURITY					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Summary: 1959-1960					
III. 3364	(CI.21E×K201)(R74×R101)	93.0	22.2	83.2	93.9
III. 3355	(H49×H51)(R71×R109B)	92.1	20.5	90.3	97.6
III. 3360	(H49×H51)(R101×Oh41)	92.0	21.0	83.4	97.0
III. 1856	(CI.21E×K201)(Oh7×38-11)	86.8	22.4	88.1	98.3
III. 1851	(Oh7×CI.21E)(38-11×C103)	78.8	20.5	93.2	98.1
Average		88.5	21.3	87.6	97.0
Number in range		Difference necessary for significance			
2-5		N.S.	N.S.	N.S.	N.S.
1960 results					
AES 904	(white) (K64×Mo22)(T111×T115)	88.7	26.8	90.6	97.7
III. 1349	(K155×K201)(38-11×Mo940)	94.4	22.4	78.3	89.3
III. 1539A	(K201×CI.21E)(38-11×CI.7)	97.4	23.1	77.2	97.7
III. 1657	(K201×CI.21E)(K4×Oh7)	106.5	24.1	76.5	100.0
III. 1660	(Oh7×CI.21E)(K4×K201)	99.3	24.2	79.2	96.2
III. 1851	(check) (Oh7×CI.21E)(38-11×C103)	95.2	21.3	89.8	96.9
III. 1856	(CI.21E×K201)(Oh7×38-11)	104.6	23.2	81.8	100.0
III. 3129	(K201×38-11)(R101×Mo01930)	106.4	20.4	80.6	96.2
III. 3133	(K201×38-11)(R127×Mo0221)	98.4	21.9	81.2	93.9
III. 3135	(K201×38-11)(R71A×Mo0221)	107.2	20.6	76.8	94.6
III. 3140	(K201×38-11)(CI.21E×Ky126)	82.4	22.6	76.9	89.3
III. 3154	(K201×CI.21E)(R132×R134)	95.1	23.6	76.8	93.9
III. 3190	(K201×C103)(Ky126×Oh7B)	95.6	22.5	81.5	87.8
III. 3193	(38-11×K12)(K201×Oh7B)	113.8	22.7	87.8	98.4
III. 3198A	(K201×Ky126)(N82481×Oh7B)	84.7	22.8	73.7	97.7
III. 3204A	(K201×Ky126)(C103×K12)	95.6	24.5	82.2	91.6
III. 3210	(CI.21E×Ky126)(C103×K12)	83.7	24.7	85.2	95.4
III. 3214	(K201×Ky126)(K12×Oh7B)	97.6	23.1	77.5	87.8
III. 3251	(K201×38-11)(K11×Ky126)	106.9	23.1	85.9	97.7
III. 3355	(H49×H51)(R71×R109B)	111.8	21.4	84.8	99.2
III. 3360	(H49×H51)(R101×Oh41)	104.7	22.2	80.4	95.4
III. 3364	(CI.21E×K201)(R74×R101)	114.7	22.2	76.3	93.1
III. 9001	(Oh7×CI.21E)(CI.7×C103)	89.8	21.5	75.8	88.6
Ind. 851	(H49×H55)(H59×B14)	102.0	22.0	73.8	93.9
Ind. 874	(H49×H52)(H59×H60)	113.3	20.7	83.9	99.2
Average		99.6	22.7	80.6	94.9
Number in range		Difference necessary for significance			
2		N.S.	1.7	N.S.	N.S.
3-5		N.S.	1.9	N.S.	N.S.
6-10		N.S.	2.0	N.S.	N.S.
11-25		N.S.	2.1	N.S.	N.S.

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (WF9 × Oh43)				
R71.....	88.9	20.4	94.2	86.8
R74.....	55.5	22.9	96.8	60.5
R76.....	88.5	21.8	91.1	84.2
R78.....	76.9	22.9	87.4	77.5
R84.....	81.5	22.4	94.6	88.3
R101.....	92.1	20.2	89.0	94.6
R104.....	78.1	20.1	95.2	79.1
R109B.....	75.9	22.6	96.9	72.9
R112.....	88.9	21.1	90.3	86.2
R113.....	73.2	20.4	93.8	77.9
R114.....	72.7	20.2	97.9	82.5
R132.....	91.9	21.6	86.8	95.4
R134.....	91.5	22.0	97.0	88.5
R151.....	83.4	22.0	96.0	75.5
R154.....	78.2	23.0	84.9	71.9
R158.....	87.1	20.5	96.3	91.3
R159.....	65.6	22.5	99.2	75.1
R166.....	79.6	21.9	81.8	81.1
R168.....	91.5	19.4	93.3	90.5
R172.....	83.8	20.5	95.6	79.5
R180.....	79.2	22.4	90.9	84.7
R181.....	92.1	18.1	91.4	87.4
R182.....	66.5	19.9	97.5	73.3
R183.....	70.7	23.3	98.4	86.8
R192.....	92.9	23.4	93.9	93.2
R193.....	83.1	21.7	89.8	86.0
R194.....	90.0	21.7	90.7	93.7
R195.....	80.2	18.2	96.8	86.5
R196.....	87.8	20.7	96.7	91.9
R197.....	91.3	24.2	90.5	84.5
R198.....	95.6	23.5	90.7	91.5
Average.....	82.4	21.4	93.1	83.8
Inbred lines crossed with (WF9 × B37)				
R71.....	95.9	23.0	96.2	90.7
R74.....	87.2	23.6	96.0	85.5
R76.....	82.1	21.6	96.4	85.8
R78.....	81.1	21.8	88.7	87.7
R84.....	68.6	21.3	96.7	85.9
R101.....	91.3	21.4	98.0	89.2
R104.....	89.3	21.0	87.3	93.8
R109B.....	75.0	23.6	94.9	82.7
R112.....	85.2	21.8	93.9	86.7
R113.....	76.0	21.1	97.2	85.1
R114.....	74.6	22.8	95.5	83.7
R132.....	85.9	21.7	81.2	82.4
R134.....	85.6	23.3	99.6	84.9
R151.....	93.8	24.8	94.7	88.7
R154.....	97.9	21.9	90.0	95.2
R158.....	61.7	21.5	96.8	67.0
R159.....	80.0	22.6	98.8	93.8
R166.....	88.8	21.5	94.6	87.4
R168.....	88.0	19.2	99.2	89.3
R172.....	84.1	22.1	99.2	87.8
R180.....	77.2	22.0	97.2	83.2
R181.....	92.4	21.2	98.7	88.8
R182.....	79.2	20.2	98.4	92.3
R183.....	64.4	24.6	99.2	80.3
R192.....	88.5	24.4	98.0	90.0
R193.....	78.3	24.2	93.0	86.5
R194.....	88.4	24.4	96.7	92.8
R195.....	80.6	21.1	96.4	85.3
R196.....	80.0	21.5	93.6	86.4
R197.....	92.7	24.1	89.0	90.8
R198.....	81.4	25.4	94.9	86.7
Average.....	83.1	22.4	95.1	87.0

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960 — continued				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (B41 × Oh7A)				
R71.....	101.6	26.3	93.7	95.0
R74.....	91.2	26.8	95.2	90.0
R76.....	85.5	26.3	92.0	90.4
R78.....	69.9	22.9	74.8	75.6
R84.....	63.6	23.7	97.9	85.1
R101.....	81.1	23.4	98.3	95.4
R104.....	86.9	23.7	94.4	82.4
R109B.....	60.2	27.1	96.7	72.0
R112.....	74.7	24.4	92.6	81.4
R113.....	70.1	23.7	97.8	88.2
R114.....	71.8	21.9	94.4	82.5
R132.....	72.9	24.9	84.4	80.6
R134.....	79.4	26.0	95.2	84.6
R151.....	85.4	24.9	93.2	81.5
R154.....	83.3	23.8	78.4	79.5
R158.....	65.1	24.5	99.2	75.6
R159.....	65.7	26.4	98.1	83.6
R166.....	93.9	25.1	79.0	97.3
R168.....	71.7	21.3	98.4	66.3
R172.....	71.8	23.7	96.6	73.4
R180.....	78.6	24.4	95.2	86.0
R181.....	76.3	22.3	88.5	73.3
R182.....	62.4	23.0	99.6	71.0
R183.....	63.1	26.7	96.3	81.5
R192.....	71.2	26.3	88.1	76.8
R193.....	76.1	24.1	94.4	82.8
R194.....	74.6	26.2	94.4	88.7
R195.....	64.4	22.9	96.4	73.3
R196.....	77.2	23.8	95.1	81.5
R197.....	77.1	28.0	94.1	83.2
R198.....	78.8	27.4	94.1	90.9
Average.....	75.8	24.7	93.1	82.2
Single-cross testers				
WF9 × Oh43.....	91.5	21.7	93.5	90.7
WF9 × B37.....	83.4	23.6	94.5	92.8
B41 × Oh7A.....	71.2	28.0	82.8	90.4
Average.....	82.0	24.4	90.3	91.3
Number in range		Difference necessary for significance		
2.....	N.S.	7.7	N.S.	11.0
3-5.....	N.S.	8.6	N.S.	12.2
6-10.....	N.S.	9.2	N.S.	13.0
11-20.....	N.S.	9.7	N.S.	13.7
21-31.....	N.S.	9.8	N.S.	13.9

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS, SUMMARY: 1959-1960 — continued				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Mean of inbred lines crossed with three testers				
R71.....	95.5	23.2	94.7	90.8
R74.....	78.0	24.4	96.0	78.7
R76.....	85.4	23.2	93.2	86.8
R78.....	76.0	22.5	83.6	80.3
R84.....	72.9	22.5	96.4	86.4
R101.....	88.2	21.7	95.0	93.1
R104.....	84.8	21.6	92.3	85.1
R109B.....	70.4	24.5	96.2	75.9
R112.....	82.9	22.4	92.3	84.8
R113.....	73.1	21.7	96.3	83.7
R114.....	73.0	21.6	95.9	82.9
R132.....	83.6	22.7	84.1	86.1
R134.....	85.5	23.8	97.3	86.0
R151.....	87.5	23.9	94.6	81.9
R154.....	86.5	22.9	84.4	82.2
R158.....	71.3	22.2	97.4	78.0
R159.....	70.4	23.9	98.7	84.2
R166.....	87.4	22.8	85.1	88.6
R168.....	83.7	20.0	97.0	82.0
R172.....	79.9	22.1	97.1	80.2
R180.....	78.3	22.9	94.4	84.6
R181.....	86.9	20.5	92.9	83.2
R182.....	69.4	21.1	98.5	78.9
R183.....	66.1	24.9	98.0	82.9
R192.....	84.2	24.7	93.3	86.7
R193.....	79.2	23.3	92.5	85.1
R194.....	84.3	24.1	93.9	91.7
R195.....	75.1	20.7	96.5	81.7
R196.....	81.7	22.0	95.1	86.6
R197.....	87.0	25.4	91.2	86.1
R198.....	85.3	25.4	93.2	89.7
Average.....	80.4	22.9	93.8	84.4

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS				
	<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (WF9 × Oh43)				
R71.....	87.4	20.8	97.1	78.0
R74.....	49.4	23.2	100.0	34.8
R76.....	90.1	22.0	98.8	81.0
R78.....	76.8	23.0	100.0	65.9
R84.....	90.8	23.1	98.3	81.8
R101.....	99.4	19.8	94.4	91.6
R104.....	76.1	19.6	100.0	65.1
R109B.....	74.1	22.2	98.7	62.9
R112.....	89.4	21.8	100.0	80.3
R113.....	76.1	20.2	100.0	65.1
R114.....	84.1	20.6	100.0	74.2
R132.....	100.1	22.2	100.0	92.4
R134.....	93.4	22.4	100.0	84.8
R151.....	71.4	21.6	100.0	59.8
R154.....	66.8	24.1	100.0	54.5
R158.....	94.1	21.2	99.1	85.6
R159.....	71.4	22.9	100.0	59.8
R166.....	80.8	21.6	100.0	70.4
R168.....	93.4	19.5	99.0	84.8
R172.....	76.8	20.3	100.0	65.9
R180.....	86.1	22.7	97.9	76.5
R181.....	89.4	17.6	100.0	80.3
R182.....	69.4	19.7	100.0	57.6
R183.....	87.4	24.5	100.0	78.0
R192.....	97.4	23.4	100.0	89.4
R193.....	87.4	22.5	100.0	78.0
R194.....	98.1	21.1	100.0	90.1
R195.....	88.1	20.8	100.0	78.8
R196.....	96.1	20.7	100.0	87.9
R197.....	88.1	25.2	94.3	78.8
R198.....	96.1	23.3	98.3	87.9
Average.....	81.8	21.7	99.2	74.9
Inbred lines crossed with (WF9 × B37)				
R71.....	96.1	22.3	100.0	87.8
R74.....	86.8	23.5	97.5	77.2
R76.....	90.1	21.1	99.0	81.0
R78.....	88.8	21.3	99.2	79.6
R84.....	86.1	21.1	99.1	76.5
R101.....	92.8	20.6	100.0	84.1
R104.....	99.4	20.2	95.2	91.6
R109B.....	86.1	22.9	98.1	76.5
R112.....	91.4	21.8	99.0	82.5
R113.....	86.1	21.1	100.0	76.5
R114.....	84.1	24.0	100.0	74.2
R132.....	82.1	21.1	96.5	71.9
R134.....	88.8	23.5	100.0	79.5
R151.....	91.4	26.1	100.0	82.5
R154.....	102.1	21.8	100.0	94.7
R158.....	60.1	22.2	96.8	47.0
R159.....	99.4	21.9	100.0	91.6
R166.....	90.1	20.6	100.0	81.0
R168.....	93.4	18.2	100.0	84.8
R172.....	89.4	21.8	100.0	80.3
R180.....	82.8	23.0	100.0	72.7
R181.....	92.1	21.8	100.0	83.3
R182.....	96.1	20.1	100.0	87.9
R183.....	80.8	25.0	100.0	70.4
R192.....	92.8	25.1	100.0	84.1
R193.....	88.8	26.0	100.0	79.5
R194.....	96.8	23.5	100.0	88.6
R195.....	88.8	21.8	100.0	79.5
R196.....	87.4	20.7	100.0	78.0
R197.....	96.8	24.2	96.5	88.6
R198.....	91.4	26.1	100.0	82.5
Average.....	89.7	21.8	99.3	80.5

(Table is continued on next page)

Table 9. — Urbana — continued

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS — continued					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Inbred lines crossed with (B41 × Oh7A)					
R71		100.1	26.1	96.2	92.4
R74		92.1	28.1	100.0	83.3
R76		92.8	27.1	100.0	84.1
R78		72.1	21.7	84.5	60.6
R84		86.1	24.7	100.0	76.5
R101		100.1	23.0	99.1	92.4
R104		87.4	24.7	100.0	78.0
R109B		67.4	27.5	100.0	55.3
R112		80.1	25.1	95.8	69.7
R113		89.4	22.9	98.3	80.3
R114		83.4	21.9	100.0	73.5
R132		80.1	25.4	100.0	69.7
R134		85.4	26.1	98.1	75.7
R151		80.8	24.6	95.5	70.4
R154		76.8	24.3	100.0	65.9
R158		72.1	25.4	100.0	60.6
R159		82.8	27.3	98.9	72.7
R166		103.4	25.5	99.2	96.2
R168		66.8	21.5	100.0	54.5
R172		70.8	24.3	100.0	59.1
R180		87.4	24.6	100.0	78.0
R181		69.4	22.5	100.0	57.6
R182		64.8	23.2	100.0	52.3
R183		81.4	28.0	97.9	71.2
R192		72.8	26.5	98.8	61.3
R193		82.1	24.6	100.0	71.9
R194		91.4	26.5	98.0	82.5
R195		68.8	23.8	100.0	56.8
R196		81.4	23.6	100.0	71.2
R197		87.4	29.2	97.3	78.0
R198		94.1	29.1	99.0	85.6
Average		82.3	25.1	98.6	72.2
Single-cross testers					
WF9 × Oh43		94.0	23.4	100.0	86.4
WF9 × B37		96.8	25.3	100.0	88.6
B41 × Oh7A		91.4	28.9	93.6	81.8
Average		94.1	25.9	97.9	85.6
Standards					
III. 1851 (C103 × 38-11)(Oh7 × CI.21E)		102.7	28.0	100.0	95.4
III. 3049 (Hy2 × WF9)(R71 × R109B)		99.3	25.4	99.2	90.9
III. 3152A (M14 × B14)(WF9 × Oh43)		100.0	20.4	99.2	93.2
III. 3347 (R74 × R101)(H49 × H55)		100.7	25.2	100.0	93.2
Average		100.7	24.8	99.6	93.2
Number in range		Difference necessary for significance			
2		14.8	11.4	4.5	16.7
3-5		16.5	12.7	5.0	18.7
6-10		17.5	13.6	5.3	19.9
11-20		18.5	14.4	5.6	21.0
21-31		19.5	15.1	6.0	22.2

(Table is concluded on next page)

Table 9. — Urbana — concluded

Entry	Acre yield	Moisture in grain	Erect plants	Stand
THREE-WAY CROSSES AND STANDARDS: 1960 RESULTS — concluded				
	<i>bu.</i>	<i>percl.</i>	<i>percl.</i>	<i>percl.</i>
Mean of inbred lines crossed with three testers				
R71.....	94.5	23.1	97.8	86.1
R74.....	76.1	24.9	99.2	65.1
R76.....	91.0	23.4	99.3	82.0
R78.....	79.2	22.0	94.6	68.7
R84.....	87.6	23.0	99.1	78.2
R101.....	97.4	21.1	97.9	89.4
R104.....	87.6	21.5	98.4	78.2
R109B.....	75.9	24.2	98.9	64.9
R112.....	87.0	22.9	98.3	77.5
R113.....	83.9	21.4	99.5	74.0
R114.....	83.9	22.2	100.0	74.0
R132.....	87.4	22.9	98.8	78.0
R134.....	89.2	24.0	99.4	80.0
R151.....	81.2	24.1	98.5	70.9
R154.....	81.9	23.4	100.0	71.7
R158.....	75.4	22.9	98.6	64.4
R159.....	84.5	24.0	99.6	74.7
R166.....	91.4	22.6	99.7	82.5
R168.....	84.5	19.7	99.7	74.7
R172.....	79.0	22.1	100.0	68.4
R180.....	85.4	23.4	99.3	75.7
R181.....	83.6	20.6	100.0	73.7
R182.....	76.8	21.0	100.0	65.9
R183.....	83.2	25.8	99.3	73.2
R192.....	87.7	25.0	99.6	78.3
R193.....	86.1	24.4	100.0	76.5
R194.....	95.4	23.7	99.4	87.1
R195.....	81.9	22.1	100.0	71.7
R196.....	88.3	21.7	100.0	79.0
R197.....	90.8	26.2	96.0	81.8
R198.....	93.9	26.2	99.1	85.3
Average.....	85.5	23.1	99.0	75.9

Table 10. — DOUBLE CROSSES OF 850 AND 900 MATURITY
Tested at Greenfield, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results, 850 maturity series					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 805	(WF9×38-11)(C103×Oh45)	89.7	20.7	97.0	75.0
Ill. 1332	(WF9×38-11)(Hy2×Oh7)	79.1	20.1	97.5	73.4
Ill. 1570	(check) (WF9×38-11)(Hy2×Oh41)	67.1	21.2	95.8	88.6
Ill. 1660	(Oh7×Cl.21E)(K4×K201)	103.0	25.3	95.4	62.8
Ill. 1976	(Oh7×Cl.21E)(38-11×Oh41)	80.3	22.9	90.6	79.5
Ill. 1978	(WF9×Oh7A)(C103×38-11)	77.6	20.8	100.0	65.9
Ill. 1996	(Hy2×Oh7)(C103×B14)	84.4	20.4	94.2	84.0
Ill. 3154	(K201×Cl.21E)(R132×R134)	88.6	27.2	88.9	61.3
Ill. 3190	(K201×C103)(Ky126×Oh7B)	102.9	26.6	96.9	74.2
Ill. 3344	(H49×H55)(R71×R105)	107.4	24.5	98.3	90.9
Ill. 3347	(H49×H55)(R74×R101)	100.9	22.3	100.0	70.4
Ill. 3348	(H49×H55)(R74×R109B)	96.6	23.8	100.0	78.7
Ill. 3350	(H49×H55)(R101×Oh41)	86.2	22.2	95.4	75.0
Ill. 3351	(H49×H55)(R109B×R168)	84.3	22.1	94.4	78.7
Ill. 3354	(H49×H51)(R71×R105)	76.4	22.3	100.0	81.0
Ill. 3357	(H49×H51)(R74×R101)	90.9	21.3	95.6	88.6
Ill. 3367	(WF9×R74)(Oh7×Cl.21E)	125.0	25.3	98.0	78.7
Ill. 3373	(WF9×C103)(R101×Oh41)	76.8	21.4	98.0	75.7
Ill. 6021	(R75×R76)(R84×K4)	71.7	21.1	98.1	71.2
Ill. 6052	(R78×38-11)(R84×K4)	79.2	21.2	90.1	76.5
Ill. 8005	(H49×H55)(B14×C103)	91.0	23.8	96.5	69.6
Ind. 851	(H49×H55)(H59×B14)	83.3	21.9	97.3	86.3
Ind. 873	(H49×H52)(H59×B14)	96.4	22.8	96.8	68.9
Ind. 874	(H49×H52)(H59×H60)	86.1	21.7	98.3	83.3
U.S. 13	(WF9×38-11)(Hy2×L317)	78.8	19.9	92.5	74.2
Average		88.1	22.5	96.2	76.5
Number in range		Difference necessary for significance			
2	20.8	2.0	6.1	N.S.
3-5	23.1	2.2	6.8	N.S.
6-10	24.5	2.4	7.2	N.S.
11-15	25.1	2.4	7.4	N.S.
16-20	25.4	2.4	7.5	N.S.
21-25	25.5	2.5	7.5	N.S.
1960 results, 900 maturity series					
AES 904 (white)	(K64×Mo22)(T111×T115)	94.0	27.3	94.5	98.4
Ill. 1349	(K155×K201)(38-11×Mo940)	90.6	21.3	93.6	96.2
Ill. 1539A	(K201×Cl.21E)(38-11×Cl.7)	75.9	24.8	87.5	84.8
Ill. 1657	(K201×Cl.21E)(K4×Oh7)	95.2	24.5	93.1	98.4
Ill. 1660	(Oh7×Cl.21E)(K4×K201)	74.9	24.6	94.4	90.1
Ill. 1851 (check)	(Oh7×Cl.21E)(38-11×C103)	94.1	23.2	94.5	96.2
Ill. 1856	(Cl.21E×K201)(Oh7×38-11)	93.9	23.7	88.1	90.1
Ill. 3129	(K201×38-11)(R101×Mo01930)	88.8	22.2	90.5	96.2
Ill. 3133	(K201×38-11)(R127×Mo0221)	81.0	22.7	88.3	85.6
Ill. 3135	(K201×38-11)(R71A×Mo0221)	65.1	22.3	93.9	87.8
Ill. 3140	(K201×38-11)(Cl.21E×Ky126)	79.4	26.7	91.3	93.9
Ill. 3154	(K201×Cl.21E)(R132×R134)	90.0	24.6	91.5	97.7
Ill. 3190	(K201×C103)(Ky126×Oh7B)	82.1	24.6	91.1	93.1
Ill. 3193	(38-11×K12)(K201×Oh7B)	77.0	24.9	91.4	90.1
Ill. 3198A	(K201×Ky126)(N82481×Oh7B)	82.6	24.4	89.5	93.9
Ill. 3204A	(K201×Ky126)(C103×K12)	83.0	26.2	91.4	96.9
Ill. 3210	(Cl.21E×Ky126)(C103×K12)	79.0	25.9	91.5	91.6
Ill. 3214	(K201×Ky126)(K12×Oh7B)	78.1	23.4	92.1	87.1
Ill. 3251	(K201×38-11)(K11×Ky126)	84.5	24.1	94.5	95.4
Ill. 3355	(H49×H51)(R71×R109B)	86.4	21.2	96.1	96.2
Ill. 3360	(H49×H51)(R101×Oh41)	93.7	21.8	98.3	92.4
Ill. 3364	(Cl.21E×K201)(R74×R101)	98.5	25.6	98.3	90.1
Ill. 9001	(Oh7×Cl.21E)(Cl.7×C103)	81.0	22.9	94.2	90.9
Ind. 851	(H49×H55)(H59×B14)	91.6	24.0	94.5	96.9
Ind. 874	(H49×H52)(H59×H60)	82.4	22.1	92.8	93.1
Average		84.9	24.0	92.7	92.9
Number in range		Difference necessary for significance			
2-25	N.S.	N.S.	N.S.	N.S.

Table 11. — DOUBLE CROSSES OF 900 MATURITY
Tested at Brownstown, 1958-1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
Summary: 1958-1960					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
Ill. 3198A	(K201×Ky126)(N82481×Oh7B)	96.6	19.3	76.6	94.0
Ill. 3190	(K201×C103)(Ky126×Oh7B)	94.7	19.7	82.4	91.8
Ill. 1660	(Oh7×CI.21E)(K4×K201)	93.1	21.1	80.3	96.1
Ill. 3193	(38-11×K12)(K201×Oh7B)	88.6	18.5	80.6	94.5
Ill. 1856	(K201×CI.21E)(Oh7×38-11)	88.6	19.8	80.5	97.9
Ill. 3214	(K201×Ky126)(K12×Oh7B)	87.4	20.2	67.9	90.7
Ill. 1851	(Oh7×CI.21E)(38-11×C103)	86.9	18.7	77.6	99.5
Ill. 1539A	(K201×CI.21E)(38-11×CI.7)	86.4	19.3	88.9	95.7
Ill. 3204A	(K201×Ky126)(C103×K12)	86.3	20.6	84.1	94.7
Ill. 3210	(CI.21E×Ky126)(C103×K12)	86.3	20.7	85.6	94.9
Ill. 3135	(K201×38-11)(R71A×Mo0221)	85.9	18.3	82.8	91.8
Ill. 3129	(K201×38-11)(R101×Mo01930)	82.8	18.7	84.8	97.0
Ill. 3133	(K201×38-11)(R127×Mo0221)	82.0	19.4	83.0	93.5
Ill. 3140	(K201×38-11)(CI.21E×Ky126)	80.7	20.3	86.2	89.4
Average		87.6	19.6	81.5	94.4
Number in range		Difference necessary for significance			
2		7.7	.8	N.S.	N.S.
3-5		8.5	.9	N.S.	N.S.
6-14		9.0	.9	N.S.	N.S.
Summary: 1959-1960					
AES 904 (white)	(K64×Mo22)(T111×T115)	83.2	23.2	45.4	95.8
Ill. 3198A	(K201×Ky126)(N82481×Oh7B)	78.1	20.8	68.1	91.7
Ill. 3154	(K201×CI.21E)(R132×R134)	75.9	22.8	63.1	92.3
Ill. 1660	(Oh7×CI.21E)(K4×K201)	74.1	22.2	72.2	95.9
Ill. 3190	(K201×C103)(Ky126×Oh7B)	73.9	20.8	73.7	88.2
Ill. 3135	(K201×38-11)(R71A×Mo0221)	69.7	19.3	73.4	87.6
Ill. 3214	(K201×Ky126)(K12×Oh7B)	69.7	21.8	52.3	85.1
Ill. 1851	(Oh7×CI.21E)(38-11×C103)	68.5	19.9	67.8	100.0
Ill. 3193	(38-11×K12)(K201×Oh7B)	68.5	20.0	71.3	93.3
Ill. 1856	(K201×CI.21E)(Oh7×38-11)	68.4	21.2	71.8	97.4
Ill. 3251	(K201×38-11)(K11×Ky126)	68.0	20.7	75.5	92.8
Ill. 1539A	(K201×CI.21E)(38-11×CI.7)	67.8	21.0	83.0	93.8
Ill. 3133	(K201×38-11)(R127×Mo0221)	66.4	20.9	77.0	89.7
Ill. 3360	(H49×H51)(R101×Oh41)	65.3	19.2	71.8	91.7
Ill. 3204A	(K201×Ky126)(C103×K12)	65.3	22.0	76.4	92.3
Ill. 3129	(K201×38-11)(R101×Mo01930)	64.9	20.2	79.7	95.8
Ill. 3210	(CI.21E×Ky126)(C103×K12)	63.5	22.2	79.4	91.8
Ill. 3355	(H49×H51)(R71×R109B)	63.1	19.9	74.4	95.3
Average		69.3	21.0	71.3	92.3
Number in range		Difference necessary for significance			
2		13.2	1.3	17.5	N.S.
3-5		14.5	1.4	19.3	N.S.
6-10		15.1	1.5	20.1	N.S.
11-19		15.4	1.5	20.4	N.S.

(Table is concluded on next page)

Table 11. — Brownstown — concluded

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 904 (white) (K64×Mo22)(T111×T115)		44.9	22.8	19.9	93.1
Ill. 1349 (K155×K201)(38-11×Mo940)		34.3	20.0	57.1	94.8
Ill. 1539A (K201×CI.21E)(38-11×CI.7)		43.3	19.8	76.7	90.5
Ill. 1657 (K201×CI.21E)(K4×Oh7)		36.8	20.6	34.1	89.7
Ill. 1660 (Oh7×CI.21E)(K4×K201)		48.1	21.9	58.8	94.0
Ill. 1851 (check) (Oh7×CI.21E)(38-11×C103)		40.1	18.7	55.5	100.0
Ill. 1856 (CI.21E×K201)(Oh7×38-11)		44.6	20.3	56.4	95.7
Ill. 3129 (K201×38-11)(R101×Mo01930)		43.3	19.3	74.5	93.1
Ill. 3133 (K201×38-11)(R127×Mo0221)		36.4	20.1	71.0	83.7
Ill. 3135 (K201×38-11)(R71A×Mo0221)		43.8	18.6	64.1	79.4
Ill. 3140 (K201×38-11)(CI.21E×Ky126)		30.0	21.2	67.4	71.7
Ill. 3154 (K201×CI.21E)(R132×R134)		49.5	22.1	50.2	87.1
Ill. 3190 (K201×C103)(Ky126×Oh7B)		42.9	19.7	61.4	82.0
Ill. 3193 (38-11×K12)(K201×Oh7B)		41.2	19.0	62.2	88.8
Ill. 3198A (K201×Ky126)(N82481×Oh7B)		51.4	19.7	62.6	86.3
Ill. 3204A (K201×Ky126)(C103×K12)		36.8	21.3	64.0	87.1
Ill. 3210 (CI.21E×Ky126)(C103×K12)		38.3	21.0	67.5	90.5
Ill. 3214 (K201×Ky126)(K12×Oh7B)		41.4	21.2	36.4	75.2
Ill. 3251 (K201×38-11)(K11×Ky126)		38.8	19.4	69.1	88.0
Ill. 3355 (H49×H51)(R71×R109B)		37.7	18.2	62.3	92.3
Ill. 3360 (H49×H51)(R101×Oh41)		37.7	17.9	63.0	86.3
Ill. 3364 (CI.21E×K201)(R74×R101)		46.7	19.3	63.7	89.7
Ill. 9001 (Oh7×CI.21E)(CI.7×C103)		43.2	19.0	57.9	90.5
Ind. 851 (H49×H55)(H59×B14)		46.3	19.4	34.5	94.8
Ind. 874 (H49×H52)(H59×H60)		40.2	20.6	63.1	90.5
Average		41.5	20.0	58.1	88.6
Number in range		Difference necessary for significance			
2		N.S.	1.8	24.9	N.S.
3-5		N.S.	2.0	27.7	N.S.
6-10		N.S.	2.1	29.3	N.S.
11-15		N.S.	2.2	30.0	N.S.
16-20		N.S.	2.2	30.5	N.S.
21-25		N.S.	2.2	30.6	N.S.

Table 12. — DOUBLE CROSSES OF 900 MATURITY
Tested at Carbondale, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 904 (white) (K64×Mo22)(T111×T115).....		103.0	28.7	100.0	72.5
Ill. 1349 (K155×K201)(38-11×Mo940).....		96.9	25.6	99.1	86.6
Ill. 1539A (K201×CI.21E)(38-11×CI.7).....		106.0	27.1	98.1	89.1
Ill. 1657 (K201×CI.21E)(K4×Oh7).....		100.6	26.9	100.0	78.3
Ill. 1660 (Oh7×CI.21E)(K4×K201).....		95.9	28.8	100.0	70.8
Ill. 1851 (check) (Oh7×CI.21E)(38-11×C103).....		104.6	26.2	100.0	80.8
Ill. 1856 (CI.21E×K201)(Oh7×38-11).....		96.3	27.4	98.3	87.5
Ill. 3129 (K201×38-11)(R101×Mo01930).....		91.8	24.6	100.0	73.3
Ill. 3133 (K201×38-11)(R127×Mo0221).....		105.5	25.7	100.0	80.0
Ill. 3135 (K201×38-11)(R71A×Mo0221).....		87.2	25.6	97.2	70.0
Ill. 3140 (K201×38-11)(CI.21E×Ky126).....		98.6	30.2	100.0	78.3
Ill. 3154 (K201×CI.21E)(R132×R134).....		111.1	27.2	98.1	90.0
Ill. 3190 (K201×C103)(Ky126×Oh7B).....		106.9	25.0	100.0	84.1
Ill. 3193 (38-11×K12)(K201×Oh7B).....		103.7	24.1	100.0	81.6
Ill. 3198A (K201×Ky126)(N82481×Oh7B).....		106.8	28.7	99.0	82.5
Ill. 3204A (K201×Ky126)(C103×K12).....		91.2	30.3	99.0	67.5
Ill. 3210 (CI.21E×Ky126)(C103×K12).....		95.1	33.1	100.0	73.3
Ill. 3214 (K201×Ky126)(K12×Oh7B).....		95.6	28.4	100.0	75.8
Ill. 3251 (K201×38-11)(K11×Ky126).....		103.2	25.5	97.2	84.1
Ill. 3355 (H49×H51)(R71×R109B).....		95.5	22.2	98.1	87.5
Ill. 3360 (H49×H51)(R101×Oh41).....		114.8	24.9	100.0	85.0
Ill. 3364 (CI.21E×K201)(R74×R101).....		98.3	26.2	100.0	80.8
Ill. 9001 (Oh7×CI.21E)(CI.7×C103).....		99.2	23.3	100.0	71.6
Ind. 851 (H49×H55)(H59×B14).....		100.6	27.3	99.1	80.0
Ind. 874 (H49×H52)(H59×H60).....		111.5	22.8	99.0	90.8
Average		100.8	26.6	99.3	80.0
Number in range		Difference necessary for significance			
2.....		N.S.	5.1	N.S.	N.S.
3-5.....		N.S.	5.6	N.S.	N.S.
6-10.....		N.S.	5.9	N.S.	N.S.
11-15.....		N.S.	6.1	N.S.	N.S.
16-25.....		N.S.	6.2	N.S.	N.S.

Table 13. — DOUBLE CROSSES OF 900 MATURITY
Tested at Wolf Lake, 1960

Entry	Pedigree	Acre yield	Moisture in grain	Erect plants	Stand
1960 results					
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>
AES 904 (white) (K64×Mo22)(T111×T115)		69.3	20.6	91.8	80.3
III. 1349 (K155×K201)(38-11×Mo940)		61.3	17.8	88.0	89.3
III. 1539A (K201×CI.21E)(38-11×CI.7)		61.4	17.1	72.1	82.5
III. 1657 (K201×CI.21E)(K4×Oh7)		64.1	18.8	84.3	87.8
III. 1660 (Oh7×CI.21E)(K4×K201)		67.2	18.6	74.8	90.9
III. 1851 (check) (Oh7×CI.21E)(38-11×C103)		59.7	17.3	90.6	90.9
III. 1856 (CI.21E×K201)(Oh7×38-11)		55.9	18.0	66.7	87.1
III. 3129 (K201×38-11)(R101×Mo01930)		78.2	17.9	87.1	91.6
III. 3133 (K201×38-11)(R127×Mo0221)		86.8	18.4	84.1	85.6
III. 3135 (K201×38-11)(R71A×Mo0221)		67.2	17.5	88.9	76.5
III. 3140 (K201×38-11)(CI.21E×Ky126)		74.7	18.5	76.9	87.8
III. 3154 (K201×CI.21E)(R132×R134)		82.3	20.3	78.4	90.9
III. 3190 (K201×C103)(Ky126×Oh7B)		66.8	17.5	83.3	88.6
III. 3193 (38-11×K12)(K201×Oh7B)		66.4	18.6	82.6	92.4
III. 3198A (K201×Ky126)(N82481×Oh7B)		66.1	18.0	70.4	83.3
III. 3204A (K201×Ky126)(C103×K12)		59.6	19.0	92.9	84.8
III. 3210 (CI.21E×Ky126)(C103×K12)		70.2	18.5	86.5	89.3
III. 3214 (K201×Ky126)(K12×Oh7B)		50.0	18.2	84.4	80.3
III. 3251 (K201×38-11)(K11×Ky126)		64.3	19.9	84.6	94.6
III. 3355 (H49×H51)(R71×R109B)		82.2	18.9	87.7	87.8
III. 3360 (H49×H51)(R101×Oh41)		73.3	18.6	76.2	93.1
III. 3364 (CI.21E×K201)(R74×R101)		81.4	19.7	83.0	85.6
III. 9001 (Oh7×CI.21E)(CI.7×C103)		61.1	17.4	76.9	91.6
Ind. 851 (H49×H55)(H59×B14)		76.9	18.6	82.7	78.0
Ind. 874 (H49×H52)(H59×H60)		70.6	17.9	70.9	99.2
Average		68.7	18.5	81.8	87.6
Number in range		Difference necessary for significance			
2	N.S.	1.8	N.S.	9.5	
3-5	N.S.	2.0	N.S.	10.5	
6-10	N.S.	2.1	N.S.	11.1	
11-15	N.S.	2.2	N.S.	11.4	
16-20	N.S.	2.2	N.S.	11.5	
21-25	N.S.	2.2	N.S.	11.6	

DOUBLE-CROSS HYBRID NUMBERS, PEDIGREES, AND INDEX TO TABLES

(The order of the single crosses does not indicate
which should be used as seed or pollen parent.)

Hybrid	Pedigree	Table No.
AES 702 (III. 1790).....	(C103 × M14) (Hy2 × WF9).....	3, 5, 6
AES 703 (III. 3019A).....	(WF9 × Oh43) (B14 × B38).....	3, 5, 6
AES 704 (III. 3016A).....	(WF9 × Oh43) (B14 × B37).....	3, 5, 6
AES 705 (III. 3011).....	(C103 × Oh43) (WF9 × B14).....	3, 4, 5, 6, 7
AES 805 (III. 1770).....	(C103 × Oh45) (WF9 × 38-11).....	8, 9, 10
AES 809.....	(C103 × Oh43) (P8 × WF9).....	4, 7
AES 810.....	(WF9 × H50) (Oh7B × Oh45).....	4, 7
AES 904W.....	(K64 × Mo22) (T111 × T115).....	9, 10, 11, 12, 13
III. 21.....	(Hy2 × 187-2) (WF9 × 38-11).....	3, 5, 6
III. 1277.....	(M14 × WF9) (I.205 × 187-2).....	2, 3, 5, 6
III. 1332.....	(Hy2 × Oh7) (WF9 × 38-11).....	8, 9, 10
III. 1349.....	(K155 × K201) (38-11 × Mo940).....	9, 10, 11, 12, 13
III. 1421.....	(WF9 × Hy2) (P8 × Oh7).....	4, 7
III. 1539A.....	(38-11 × Cl.7) (K201 × Cl.21E).....	9, 10, 11, 12, 13
III. 1555A.....	(WF9 × Oh51A) (I.224 × Oh28).....	2, 3
III. 1559B.....	(M14 × Oh28) (WF9 × Oh51A).....	2, 3
III. 1570.....	(Hy2 × Oh41) (WF9 × 38-11).....	8, 9, 10
III. 1657.....	(K201 × Cl.21E) (K4 × Oh7).....	9, 10, 11, 12, 13
III. 1660.....	(K4 × K201) (Oh7 × Cl.21E).....	8, 9, 10, 11, 12, 13
III. 1851.....	(C103 × 38-11) (Oh7 × Cl.21E).....	9, 10, 11, 12, 13
III. 1856.....	(38-11 × Oh7) (K201 × Cl.21E).....	9, 10, 11, 12, 13
III. 1861.....	(WF9 × M14) (I.224 × Oh28).....	2, 3
III. 1863.....	(WF9 × M14) (I.205 × Oh43).....	2, 3
III. 1922.....	(Hy2 × WF9) (R71 × R105).....	3, 5, 6
III. 1936.....	(Hy2 × WF9) (M14 × B14).....	2, 3, 5, 6
III. 1952.....	(M14 × B14) (A545 × W64A).....	2, 3
III. 1955.....	(M14 × A297) (B14 × W64A).....	2, 3
III. 1957.....	(M14 × A545) (B14 × W64A).....	2, 3
III. 1958.....	(M14 × Oh26A) (B14 × A545).....	2, 3
III. 1959.....	(M14 × W64A) (B14 × A297).....	2, 3
III. 1960.....	(M14 × W64A) (B14 × A545).....	2, 3
III. 1961.....	(B14 × A545) (A239 × W64A).....	2, 3
III. 1962.....	(B14 × A545) (A297 × W64A).....	2, 3
III. 1968.....	(R163 × R169) (WF9 × B14).....	3, 5, 6
III. 1969.....	(R165 × R168) (WF9 × B14).....	3, 5, 6
III. 1969A.....	(R165 × WF9) (R168 × B14).....	2, 3
III. 1976.....	(38-11 × Oh41) (Oh7 × Cl.21E).....	8, 9, 10
III. 1978.....	(C103 × 38-11) (WF9 × Oh7A).....	8, 9, 10
III. 1983.....	(Hy2 × B14) (WF9 × 38-11).....	4, 7
III. 1996.....	(C103 × B14) (Hy2 × Oh7).....	4, 7, 8, 9, 10
III. 3009.....	(B14 × B21) (A297 × W64A).....	2, 3
III. 3022.....	(WF9 × B14) (N22A × Oh43).....	3, 5, 6
III. 3029.....	(WF9 × B14) (Oh43 × Oh45).....	3, 5, 6
III. 3042.....	(WF9 × B14) (B40 × Oh45).....	3, 4, 5, 6, 7
III. 3049.....	(Hy2 × WF9) (R71 × R109B).....	4, 7
III. 3080.....	(Hy2 × WF9) (R101 × Oh451).....	4, 7
III. 3129.....	(R101 × Mo01930) (38-11 × K201).....	9, 10, 11, 12, 13
III. 3133.....	(R127 × Mo0221) (38-11 × K201).....	9, 10, 11, 12, 13
III. 3135.....	(R71A × Mo0221) (38-11 × K201).....	9, 10, 11, 12, 13
III. 3140.....	(38-11 × K201) (Ky126 × Cl.21E).....	9, 10, 11, 12, 13

(Index is concluded on next page)

Index to tables — concluded

Hybrid	Pedigree	Table No.
III. 3152	(M14 × WF9) (B14 × Oh43)	2, 3, 5, 6
III. 3154	(R132 × R134) (K201C × Cl.21E)	8, 9, 10, 11, 12, 13
III. 3160	(WF9 × Oh7) (B14 × Oh43)	4, 7
III. 3173	(B14 × Oh43) (A545 × N24)	2, 3
III. 3174	(B37 × Oh28) (A297 × Oh43)	2, 3
III. 3182A	(R105 × WF9) (R151 × R154)	3, 5, 6
III. 3183	(R105 × R153) (R154 × WF9)	4, 7
III. 3190	(C103 × K201) (Ky126 × Oh7B)	8, 9, 10, 11, 12, 13
III. 3193	(38-11 × K12) (K201 × Oh7B)	9, 10, 11, 12, 13
III. 3198A	(N82481 × Oh7B) (K201 × Ky126)	9, 10, 11, 12, 13
III. 3204A	(C103 × K12) (K201 × Ky126)	9, 10, 11, 12, 13
III. 3210	(C103 × K12) (Ky126 × Cl.21E)	9, 10, 11, 12, 13
III. 3214	(K201 × Ky126) (K12 × Oh7B)	9, 10, 11, 12, 13
III. 3237	(R101 × WF9) (R151 × R154)	4, 7
III. 3244	(R105 × R153) (R151 × WF9)	4, 7
III. 3251	(38-11 × K201) (K11 × Ky126)	9, 10, 11, 12, 13
III. 3265	(R71 × R109B) (WF9 × Oh43)	3, 5, 6
III. 3266	(R74 × R109B) (WF9 × Oh43)	3, 5, 6
III. 3270	(R74 × R168) (WF9 × Oh43)	3, 5, 6
III. 3275	(R114 × R168) (WF9 × Oh43)	3, 5, 6
III. 3291	(P8 × WF9) (B14 × Oh43)	4, 7
III. 3294	(C103 × Hy2) (P8 × WF9)	4, 7
III. 3301	(M14 × Oh43) (R168 × B14)	2, 3
III. 3302A1	(M14 × W64A) (R172 × B14)	2, 3
III. 3303	(M14 × Oh43) (R172 × B14)	3, 5, 6
III. 3313	(L12 × B14) (Oh43 × W64A)	2, 3
III. 3315A	(Hy2 × WF9) (R109B × B14)	3, 5, 6
III. 3343	(R71 × R74) (H49 × H55)	4, 7
III. 3344	(R71 × R105) (H49 × H55)	8, 9, 10
III. 3346	(R71 × R168) (H49 × H55)	4, 7
III. 3347	(R74 × R101) (H49 × H55)	3, 5, 6, 8, 9, 10
III. 3348	(R74 × R109B) (H49 × H55)	4, 7, 8, 9, 10
III. 3350	(R101 × Oh41) (H49 × H55)	8, 9, 10
III. 3351	(R109B × R168) (H49 × H55)	4, 7, 8, 9, 10
III. 3354	(R71 × R105) (H49 × H51)	8, 9, 10
III. 3355	(R71 × R109B) (H49 × H51)	9, 10, 11, 12, 13
III. 3357	(R74 × R101) (H49 × H51)	8, 9, 10
III. 3360	(R101 × Oh41) (H49 × H51)	9, 10, 11, 12, 13
III. 3364	(R74 × R101) (K201 × Cl.21E)	9, 10, 11, 12, 13
III. 3367	(R74 × WF9) (Oh7 × Cl.21E)	8, 9, 10
III. 3373	(C103 × WF9) (R101 × Oh41)	8, 9, 10
III. 3381	(R71 × WF9) (B14 × Oh43)	3, 5, 6
III. 3382	(R109B × WF9) (B14 × Oh43)	3, 5, 6
III. 3383	(M14 × WF9) (R172 × Oh43)	3, 5, 6
III. 3384	(Hy2 × Oh7) (WF9 × Oh41)	4, 7
III. 6021	(R75 × R76) (R84 × K4)	8, 9, 10
III. 6052	(R78 × 38-11) (R84 × K4)	8, 9, 10
III. 6201	(R53 × Oh7) (WF9 × B14)	2, 3
III. 6202	(R53 × Oh51) (Oh43 × W64A)	2, 3
III. 8001	(Hy2 × R138) (Oh7 × Oh7B)	4, 7
III. 8002	[(Hy2 × B14) Hy2] [(Oh7 × C103) Oh7]	4, 7
III. 8003	(WF9 × Oh7) (H55 × C103)	4, 7
III. 8004	(WF9 × Hy2) (R74 × B14)	4, 7
III. 8005	(H49 × H55) (B14 × C103)	8, 9, 10
III. 9001	(Oh7 × Cl.21E) (Cl.7 × C103)	9, 10, 11, 12, 13
Ind. 851	(H49 × H55) (H59 × B14)	8, 9, 10, 11, 12, 13
Ind. 873	(H49 × H52) (H59 × B14)	8, 9, 10
Ind. 874	(H49 × H52) (H59 × H60)	8, 9, 10, 11, 12, 13
U.S. 13	(WF9 × 38-11) (Hy2 × L317)	4, 7, 8, 9, 10





UNIVERSITY OF ILLINOIS-URBANA

Q.630.71L68
BULLETIN. URBANA
869 1981

C008



3 0112 019530341